

REVISED VERSION

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 July 2004 (08.07.2004)

PCT

(10) International Publication Number
WO 2004/056866 A1

(51) International Patent Classification⁷: C07K 14/705,
A61K 38/17, A61P 11/06, C12N 5/12, C12Q 1/68

(74) Agent: OY JALO ANT-WUORINEN AB; Iso
Roobertinkatu 4-6 A, FIN-00120 HELSINKI (FI).

(21) International Application Number:
PCT/FI2003/000973

(22) International Filing Date:
19 December 2003 (19.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/435,846 20 December 2002 (20.12.2002) US
60/437,895 3 January 2003 (03.01.2003) US
60/458,767 26 March 2003 (26.03.2003) US
60/486,000 9 July 2003 (09.07.2003) US

(71) Applicant (for all designated States except US): GENEOS
OY [FI/FI]; Tukholmankatu 2, 2. krs, FI-00250 Helsinki
(FI).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LAITINEN, Tarja
[FI/FI]; Myllytie 18, FIN-04400 JÄRVENPÄÄ (FI).
KERE, Juha [FI/FI]; Lahntie 1, FIN-02170 ESPOO
(FI). LAITINEN, Lauri, A. [FI/FI]; Katiskakuja 3 as 6,
FIN-02230 ESPOO (FI). POLVI, Anne [FI/FI]; Kalliop-
urontie 21 A 4, FIN-02920 ESPOO (FI). MÄKELÄ,
Siru [FI/FI]; Sankaritie 7 A 12, FIN-00320 HELSINKI
(FI). VENDELIN, Johanna [FI/FI]; Kilonrinne 7 B8,
FIN-02610 ESPOO (FI). PULKKINEN, Ville [FI/FI];
Strömbergintie 14 B 41, FIN-00380 HELSINKI (FI).
SALMIKANGAS, Paula [FI/FI]; Aurankaari 2 B 8,
FIN-01800 KLAUKKALA (FI).

(81) Designated States (*national*): AE, AG, AL, AM, AT (util-
ity model), AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,
CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE
(utility model), DE, DK (utility model), DK, DM, DZ, EC,
EE (utility model), EE, EG, ES, FI (utility model), FI, GB,
GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG,
KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL,
PT (utility model), PT, RO, RU, SC, SD, SE, SG, SK (util-
ity model), SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG,
US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the revised international search
report: 21 July 2005

(15) Information about Correction:
see PCT Gazette No. 29/2005 of 21 July 2005, Section II

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: ASTHMA SUSCEPTIBILITY LOCUS

(57) Abstract: The present invention describes a susceptibility locus which is functionally related to asthma. The locus maps within human chromosome 7p15-p14. The invention also describes a novel human gene, GPRA. The invention provides diagnostic methods and materials for analysing allelic variation in said locus and the GPRA gene. The invention also provides polypeptides encoded by GPRA gene and antibodies binding to said polypeptides. The invention further provides pharmaceutical compositions for the treatment of asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer.

WO 2004/056866 A1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 2003/000973

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07K 14/705, A61K 38/17, A61P 11/06, C12N 5/12, C12Q 1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, BIOSIS, MEDLINE, CHEM.ABS.DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 0148188 A1 (HELIX RESEARCH INSTITUTE); 5 July 2001 (05.07.2001), abstract, Figure 11, & CAS RN 349590-17-6, & GenBank accession no. B0095689 --	1-3,5-17, 19-23,25, 27-28,32-44, 47-62,64,
X	WO 0118206 A1 (LEXICON GENETICS INCORPORATED), 15 March 2001 (15.03.2001), see the entire document, especially SEQ.ID.Nos. 1-2, p.5, 1.30 - p.6, 1.19, p.6, 1.25 - p.7, 1.1, p.7, 1.26 - p.8, 1.7, p.14, 1.23-32, p.10, 1.19-30 --	1-3,5-17, 19-23,25, 27-28,32-44, 47-62,64,
X	WO 0242458 A2 (TULARIK INC.), 30 May 2002 (30.05.2002), see the entire document, especially pages 75-76, [58], [71], p. 83, claim 33, [217], lines 32-33, [49] --	1-24,32-33, 56-62,66-69, 110,112,115

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

6 April 2005

Date of mailing of the international search report

14 -04- 2005

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

SARA NILSSON/EÖ

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 2003/000973

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 0231145 A1 (TAKEDA CHEMICAL INDUSTRIES, LTD.), 18 April 2002 (18.04.2002), SEQ.ID.No. 16, & CAS RN 413009- 26-4 --	1-3,5-17, 19-23,27-28, 32-44,47-62, 64,
X	WO 0148015 A2 (PHARMACIA & UPJOHN COMPANY), 5 July 2001 (05.07.2001), SEQ.ID.No. 25 and 12, p.10, 1.9-12, p.14, 1.4-6, p.16, 1.15-26, p.19, 1.21-24. p.62, 1.1-18, p.69, 1.17 - p. 71, 1. 24 --	1-25,27-28, 32-44,47-62, 64,66-69, 110,112,
X	WO 0177137 A1 (HUMAN GENOME SCIENCE, INC.), 18 October 2001 (18.10.2001), SEQ.ID.No. 1962, & CAS RN 369415-18-9 --	1-3,5,8-14, 110
X	DATABASE WPI Week 200170 Derwent Publication Ltd., London, GB; Class B04, AN 2001-610074 & JP 2001245666 A (KYOWA HAKKO KOGYO KK), 11 September 2001 (2001-09-11) abstract, SEQ.ID.No. 19, & GenBank acc.no. B0017045 --	1-3,5-17, 19-23,25, 27-28,32-44, 47-62,64,
X	WO 02063004 A2 (INCYTE GENOMICS, INC.), 15 August 2002 (15.08.2002), SEQ.ID.No. 51, p. 28, 1.2-13, p. 77, 1.14-27, p. 78, 1.6-12, p. 44, 1.14-32 --	1-25,27-28, 32-44,47-62, 64,66-69, 110,112,
X	WO 0196400 A2 (MILLENNIUM PHARMACEUTICALS, INC.), 20 December 2001 (20.12.2001), SEQ.ID.No. 1 and 2, p. 70, 1. 1-21, p. 69, 1. 6-22 --	1-3,5-17, 19-23,25, 27-28,32-44, 47-62,
X	WO 0218412 A1 (HUMAN GENOMICS SCIENCES, INC.), 7 March 2002 (07.03.2002), SEQ.ID.No. 53, & CAS RN 402537-39-7 and 402537-11-5 --	1-3,5,8-14, 110

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 2003/000973

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	European Journal of Human Genomics, Volume 10, 2002, Anne Polvi et al, "Physical map of an asthma susceptibility locus in 7p15-p14 and an association study of TCRG", pages 658-665, see entrie article --	1-119
X	Database EMBL Acc. no. BC031961 28-Jun-2992 & Strausberg et al: PNAS, December 24, 2002, vol. 99, no. 26, pages 16899-16903 --	70-85,89-90, 104-108
P,X	WO 03007187 A1 (TAKEDA CHEMICAL INDUSTRIES, LTD.), 23 January 2003 (23.01.2003), SEQ.ID.No. 37, figures15-17, & CAS RN 349588-91-6 --	1-3,5,8-17, 19-23,56-59, 66-69
P,X	WO 03025179 A1 (TAKEDA CHEMICAL INDUSTRIES, LTD.), 27 March 2003 (27.03.2003), SEQ.ID.No. 67, figures 41-42, & CAS RN 503077-41-6 --	1-3,5,8-17, 19-23,27-28, 32-45,47-62, 64
P,X	EP 1365030 A1 (MTM LABORATORIES AG), 26 November 2003 (26.11.2003), SEQ.ID.No. 1 (0021), abstract, figure 5 -- -----	1-25,27-28, 32-44,47-62, 64,66-69, 110,112

INTERNATIONAL SEARCH REPORT

International application No.
PCT/FI2003/000973

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **25-31 and 34-46 partially**
because they relate to subject matter not required to be searched by this Authority, namely:
see extra sheet
2. ☐ Claims Nos.: **5, 15 and 78**
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
see extra sheet
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

<Box II.1>

Claims 25-31 and 86-88 relate to methods of treatment of the human or animal body by surgery or by therapy or diagnostic methods practiced on the human or animal body (PCT Rule 39.1(iv)). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds or compositions and on the polypeptides per se.

Claims 34-46 and 91-96 may relate to methods of treatment of the human or animal body by surgery or by therapy or diagnostic methods practiced on the human or animal body (PCT Rule 39.1(iv)) since it is not clear whether the method is performed in vitro or in vivo. Nevertheless, a search has been executed for these claims. The search has been based on the assumption that the method is performed entirely in vitro.

<Box II.2>

Present claims 5 and 15 relate to an extremely large number of possible polypeptides. Support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the polypeptides claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Consequently, the search has been focused on those parts of the claims which appear to be supported and disclosed, namely those parts related to the sequences SEQ ID NOs: 3, 5, 7, 9, 11, 13 and 15 and related sequences including the variants shown in Table 7.

Present claim 78 relate to an extremely large number of possible nucleic acids. Support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the nucleic acids claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts related to the nucleic acids SEQ ID NO:s 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, and 40.

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/03/2005

International application No.

PCT/FI 2003/000973

WO	0148188	A1	05/07/2001	AU	2230301 A	09/07/2001
				AU	2230401 A	09/07/2001
				EP	1243648 A	25/09/2002
				US	20030157558 A	21/08/2003
				WO	0148189 A	05/07/2001
				AU	4467301 A	08/10/2001
				WO	0173023 A	04/10/2001

WO	0118206	A1	15/03/2001	AU	7475400 A	10/04/2001
				CA	2384663 A	15/03/2001
				EP	1254220 A	06/11/2002
				JP	2003508076 T	04/03/2003
				US	20030130495 A	10/07/2003

WO	0242458	A2	30/05/2002	AU	3929102 A	03/06/2002
				US	20030027252 A	06/02/2003

WO	0231145	A1	18/04/2002	AU	9423901 A	22/04/2002
				EP	1329508 A	23/07/2003
				JP	2002355061 A	10/12/2002
				US	20040110920 A	10/06/2004

WO	0148015	A2	05/07/2001	AU	5787501 A	09/07/2001
				CA	2394749 A	05/07/2001
				EP	1242454 A	25/09/2002
				US	20020062013 A	23/05/2002
				AU	4165801 A	03/09/2001
				EP	1265925 A	18/12/2002
				US	20030003451 A	02/01/2003
				WO	0162797 A	30/08/2001
				AU	4165501 A	03/09/2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/03/2005

International application No.

PCT/FI 2003/000973

WO	0177137	A1	18/10/2001	AU	5906301 A	30/10/2001
				AU	5906601 A	30/10/2001
				AU	6102401 A	30/10/2001
				AU	6294201 A	30/10/2001
				AU	6456301 A	30/10/2001
				AU	6655701 A	23/10/2001
				AU	7480901 A	30/10/2001
				CA	2405525 A	25/10/2001
				CA	2405550 A	25/10/2001
				CA	2405557 A	25/10/2001
				CA	2405563 A	25/10/2001
				CA	2405701 A	25/10/2001
				CA	2405709 A	25/10/2001
				CA	2405912 A	18/10/2001
				EP	1274719 A	15/01/2003
				EP	1274720 A	15/01/2003
				EP	1276756 A	22/01/2003
				EP	1276849 A	22/01/2003
				EP	1276856 A	22/01/2003
				EP	1278544 A	29/01/2003
				EP	1278767 A	29/01/2003
				JP	2003530838 T	21/10/2003
				JP	2003530839 T	21/10/2003
				JP	2003530846 T	21/10/2003
				JP	2003530847 T	21/10/2003
				JP	2003530852 T	21/10/2003
				JP	2003531590 T	28/10/2003
				JP	2004506407 T	04/03/2004
				US	20030125247 A	03/07/2003
				US	20030171267 A	11/09/2003
				US	20030199043 A	23/10/2003
				US	20030219875 A	27/11/2003
				US	20040010134 A	15/01/2004
				WO	0179258 A	25/10/2001
				WO	0179271 A	25/10/2001
				WO	0179442 A	25/10/2001
				WO	0179443 A	25/10/2001
				WO	0179444 A	25/10/2001
				WO	0179480 A,B	25/10/2001

WO	02063004	A2	15/08/2002	CA	2437717 A	15/08/2002
				EP	1366165 A	03/12/2003

WO	0196400	A2	20/12/2001	AU	6853101 A	24/12/2001
				EP	1294766 A	26/03/2003
				US	20020151046 A	17/10/2002

WO	0218412	A1	07/03/2002	AU	2950301 A	13/03/2002
				CA	2420392 A	07/03/2002
				EP	1317472 A	11/06/2003

WO	03007187	A1	23/01/2003	JP	2003159095 A	03/06/2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/03/2005

International application No.

PCT/FI 2003/000973

WO	03025179	A1	27/03/2003	CA	2460468 A	27/03/2003
				EP	1433849 A	30/06/2004
				JP	2004065232 A	04/03/2004

EP	1365030	A1	26/11/2003	WO	03097872 A	27/11/2003

Figs. 12A and 12B: In Western blot analysis of several human tissues, the A variant specific antibody recognized four major polypeptide bands at approximately 40, 42, 44, and 50 kDa (12A). 50 kDa band was detectable in skeletal muscle whereas uterine muscle, colon epithelium and prostate showed similar expression patterns of *GPRA A* at approximately 50, 44, 40 kDa. The most intensive *GPRA A* expression was recorded in colon muscle at 42 kDa. The B variant specific antibody recognized two major polypeptide bands at approximately 25 and 39 kDa with the most intensive *GPRA* expression in kidney (12B).

Fig. 13: Genomic structure of AAA1 showing both the exon (size below) and intron (size above) structures of the gene. AAA1 (asthma associated alternatively spliced gene 1) shows complex splicing, two alternative starting methionines and six alternative stop codons for the predicted polypeptide was identified. Location of the primers used in cloning of different splice variants are shown below. Gray area shows the location of AST1.

Fig. 14: I-XII different splice variants of AAA1, I-XI are full length cDNAs.

Fig. 15: Sequence alignments of the predicted polypeptides encoded by the I-XI splice variants of AAA1 (CLUSTAL W program). The conserved region of the AAA1 encoded polypeptides is in bold.

Fig. 16: Human multiple tissue expression array (BD, Clontech) hybridized with the probe specific for AAA1 (a mixture of multiple splice variants) (upper panel) and location of studied RNAs in the dot blot (lower panel).

Fig. 17: Human fetal multiple tissue northern blot (BD, Clontech) hybridized with the probe specific for AAA1 (a mixture of multiple splice variants) identifies several alternative transcripts (arrows).

Fig. 18: Tissue specific expression AAA1. RT-PCR with the variants I, IV, VI, X, and XI specific primer pairs in liver (Li), lung (Lu), testis (Te), and kidney (Ki) shows different patterns of transcripts in different tissues.

Fig. 19: Variable alternative splicing for AAA1 depending on genotype. RT-PCR spanning exons 6 to 10b of AAA1 was performed on lymphoblastic RNA samples genotyped for AST1. Only the non-carrier of AST1 processes normal amount of the exon 6-10b transcript, whereas a homozygote and heterozygotes show either absent transcript or smaller splice variants. Beta-actin was used as control in parallel amplifications.

Fig. 20: Nucleic (SEQ ID NOS:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40) and amino acid sequences (SEQ ID NOS:17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41) of splice variants of AAA1.

Figs. 21A and 21B: 21A. *In vitro* translation of AAA1 gene using rabbit reticulocyte lysate translation machinery with S³⁵-labelled methionine in the reaction mixture. 21B. Specificity of the AAA1 antibody (anti-AAA1). Glutathione S-transferase (GST) -fusion proteins for AAA1 were expressed in *E. coli* as GST fusion proteins (GST-AAA1).

DEFINITIONS

Unless defined otherwise, all technical and scientific terms have the same meaning as is commonly understood by one of skill in the art to which this invention is related. The definitions below are presented for clarity.

"Isolated" when referred to a molecule, refers to a molecule that has been identified and separated and/or recovered from a component of its natural environment and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide. The term "isolated" does not refer to genomic or cDNA libraries, whole cell total or mRNA preparations, genomic DNA preparations (including those separated by electrophoresis and transferred onto blots), sheared whole cell genomic DNA preparations or other compositions where the art demonstrates no distinguishing features of the polynucleotide sequences of the present invention.

"Nucleic acid", includes DNA molecules (*e.g.* cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), analogs of the DNA or RNA generated using nucleotide analogs, and derivatives, fragments and homologs. The nucleic acid or nucleic acid molecule may be single-stranded or double-stranded, but preferably comprises double-stranded DNA. Preferred nucleic acids of the invention include segments of DNA, or their complements including any one of the polymorphic sites shown in Table 3, 7 or 12. The segments are usually between 5 and 100 contiguous bases, and often range from 5, 10, 12, 15, 20, or 25 nucleotides to 10, 15, 30, 25, 20, 50 or 100 nucleotides. Nucleic acids between 5-10, 5-20, 10-20, 12-30, 15-30, 10-50, 20-50 or 20-100 bases are common. The polymorphic site can occur within any position of the segment. The segments can be from any of the allelic forms of DNA shown in Table 3, 7 or 12. For brevity in Table 3, the symbol T is used to represent both thymidine in DNA and uracil in RNA. Thus, in RNA oligonucleotides, the symbol T should be construed to indicate a uracil residue. Unless otherwise apparent from the context,

reference to a SEQ ID NO. of the invention, refers to the strand shown in the sequence listing for that SEQ ID NO., the perfect complementary strand thereof or a duplex of the two. Thus, for example, reference to primers hybridizing to SEQ ID NO:1 includes primers hybridizing to the strand shown in the sequence listing and the complementary strand.

"Isolated nucleic acid" is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an isolated nucleic acid is free of sequences that naturally flank the nucleic acid (*i.e.* sequences located at the 5'- and 3'-termini of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, isolated asthma locus-1 molecules can contain less than about 5 kb, 4 kb, 3kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell/tissue from which the nucleic acid is derived (*e.g.*, brain, heart, liver, spleen, etc.). Moreover, an isolated nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material or culture medium when produced by recombinant techniques, or of chemical precursors or other chemicals when chemically synthesized.

The term "fragment" or "segment" as applied to a longer nucleic acid, may ordinarily be at least about 10 contiguous nucleotides of the longer nucleic acid in length, typically, at least about 20 contiguous nucleotides, more typically, from about 20 to about 50 contiguous nucleotides, preferably at least about 50 to about 100 contiguous nucleotides, even more preferably at least about 100 contiguous nucleotides to about 300 contiguous nucleotides, yet even more preferably at least about 300 to about 400, and most preferably, the nucleic acid fragment will be greater than about 500 contiguous nucleotides in length.

"Oligonucleotide" comprises a series of linked nucleotide residues, which oligonucleotide has a sufficient number of nucleotide bases to be used in a PCR reaction or other application. A short oligonucleotide sequence may be based on, or designed from, a genomic or cDNA sequence and is used to amplify, confirm, or reveal the presence of an identical, similar or complementary DNA or RNA in a particular cell or tissue. Oligonucleotides comprise portions of a nucleic acid.

"Variant" refers to a polynucleotide differing from the polynucleotide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely

similar, and, in many regions, identical to the polynucleotide of the present invention. Allelic variants of a gene refer to variant forms of the same gene between different individuals of the same species. The first identified allelic form is arbitrarily designated as the reference form and other allelic forms are designated as alternative or variant alleles. The allelic form occurring most frequently in a selected population is sometimes referred to as the wildtype form. Diploid organisms may be homozygous or heterozygous for allelic forms. A diallelic polymorphism has two forms. A triallelic polymorphism has three forms. Cognate forms of a gene refers to variation between structurally and functionally related genes between species. For example, the human gene showing the greatest sequence identity and closest functional relationship to a mouse gene is the human cognate form of the mouse gene. Thus, for example, the invention includes primate, bovine, ovine, murine, and avian cognate forms of the human GPRA and AAA1 genes.

A single nucleotide polymorphism occurs at a polymorphic site occupied by a single nucleotide, which is the site of variation between allelic sequences. The site is usually preceded by and followed by highly conserved sequences of the allele (*e.g.*, sequences that vary in less than 1/100 or 1/1000 members of the populations). A single nucleotide polymorphism usually arises due to substitution of one nucleotide for another at the polymorphic site. A transition is the replacement of one purine by another purine or one pyrimidine by another pyrimidine. A transversion is the replacement of a purine by a pyrimidine or vice versa. Single nucleotide polymorphisms can also arise from a deletion of a nucleotide or an insertion of a nucleotide relative to a reference allele.

A set of polymorphisms means at least 2, and sometimes 5, or more of the polymorphisms shown in Table 3, 7 or 12.

"Stringency". Highly stringency conditions are well-known in the art, *e.g.* 6×NaCl/sodium citrate (SSC) at about 45 °C is applied for a hybridization step, followed by a wash of 2×SSC at 50 °C or , *e.g.*, alternatively hybridization at 42 °C in 5×SSC, 20 mM NaPO₄, pH 6.8, 50% formamide; and washing at 42 °C in 0.2×SSC. These conditions can be varied empirically based on the length and the GC nucleotide base content of the sequences to be hybridized, or based on formulas for determining such variation (See, for example, Sambrook *et al.*, "Molecular Cloning: A Laboratory Manual", Second Edition, pages 9.47-9.51, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1989)). A low stringency is

defined herein as being in 4-6 X SSC/0.1-0.5% w/v SDS at 37-45 degree of C for 2-3 hours. Depending on the source and concentration of nucleic acid involved in the hybridization, alternative conditions of stringency may be employed such as medium stringent conditions which are considered herein to be 1-4 X SSC/0.25-0.5% w/v SDS at 45 degree of Celsius for 2-3 hours or highly stringent conditions considered herein to be 0.1-1 X SSC/0.1% w/v SDS at 60 degree of Celsius for 1-3 hours.

Hybridization probes are capable of binding in a base-specific manner to a complementary strand of nucleic acid. Such probes include nucleic acids, peptide nucleic acids, as described in Nielsen *et al.*, Science 254, 1497-1500 (1991). "Probes" are nucleic acid sequences of variable length, preferably between at least about 10 nucleotides (nt), 100 nt, or many (*e.g.*, 6000 nt) depending on the specific use. Often probes have 15-50 nucleotides. Probes are used to detect identical, similar, or complementary nucleic acid sequences. Longer length probes can be obtained from a natural or recombinant source, are highly specific, and much slower to hybridize than shorter-length oligomer probes. Probes may be single- or double-stranded and designed to have specificity in PCR, membrane-based hybridization technologies, or ELISA-like technologies. Probes also hybridize to nucleic acid molecules in biological samples, thereby enabling immediate applications in chromosome mapping, linkage analysis, tissue identification and/or typing, and a variety of forensic and diagnostic methods of the invention.

The term primer refers to a single-stranded oligonucleotide capable of acting as a point of initiation of template-directed DNA synthesis under appropriate conditions (*i.e.*, in the presence of four different nucleoside triphosphates and an agent for polymerization, such as, DNA or RNA polymerase or reverse transcriptase) in an appropriate buffer and at a suitable temperature. The appropriate length of a primer depends on the intended use of the primer but typically ranges from 15 to 30, 40 or 50 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template but must be sufficiently complementary to hybridize with a template. The term primer site refers to the area of the target DNA to which a primer hybridizes. The term primer pair means a set of primers including a 5' upstream primer that hybridizes with the 5' end of the DNA sequence to be amplified and a 3', downstream primer that hybridizes with the complement of the 3' end of the sequence to be amplified.

"Vector" means any plasmid or virus encoding an exogenous nucleic acid. The term should also be construed to include non-plasmid and non-viral compounds which facilitate transfer of nucleic acid into virions or cells, such as, for example, polylysine compounds and the like. The vector may be a viral vector which is suitable as a delivery vehicle for delivery of the nucleic acid encoding the desired protein, or mutant thereof, to a cell, or the vector may be a non-viral vector which is suitable for the same purpose. Examples of viral and non-viral vectors for delivery of DNA to cells and tissues are well known in the art and are described, for example, in Ma *et al.* (1997, Proc. Natl. Acad. Sci. U.S.A. 94:12744-12746). Examples of viral vectors include, but are not limited to, a recombinant vaccinia virus, a recombinant adenovirus, a recombinant retrovirus, a recombinant adeno-associated virus, a recombinant avian pox virus, and the like (Cranage *et al.*, 1986, EMBO J. 5:3057-3063; International Patent Application No. W094/17810, published August 18, 1994; International Patent Application No. W094/23744, published October 27, 1994). Examples of non-viral vectors include, but are not limited to, liposomes, polyamine derivatives of DNA, and the like.

"Recombinant polynucleotide" refers to a polynucleotide having sequences that are not naturally joined together. An amplified or assembled recombinant polynucleotide may be included in a suitable vector, and the vector can be used to transform a suitable host cell.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, or cows. Preferably, the mammal is human.

Linkage describes the tendency of genes, alleles, loci or genetic markers to be inherited together as a result of their location on the same chromosome, and can be measured by percent recombination between the two genes, alleles, loci or genetic markers that are physically-linked on the same chromosome. Loci occurring within 50 centimorgan of each other are linked. Some linked markers occur within the same gene or gene cluster.

Polymorphism refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. A polymorphic marker or site is the locus at which divergence occurs. Preferred markers have at least two alleles, each occurring at frequency

of greater than 1%, and more preferably greater than 10% or 20% of a selected population. A polymorphic locus may be as small as one base pair.

Linkage disequilibrium (LD) or allelic association means the preferential association of a particular allele or genetic marker with a specific allele, or genetic marker at a nearby chromosomal location more frequently than expected by chance for any particular allele frequency in the population. For example, if locus X has alleles a and b, which occur equally frequently, and linked locus Y has alleles c and d, which occur equally frequently, one would expect the haplotype ac to occur with a frequency of 0.25 in a population of individuals. If ac occurs more frequently, then alleles a and c are considered in linkage disequilibrium.

Linkage disequilibrium may result from natural selection of certain combination of alleles or because an allele has been introduced into a population too recently to have reached equilibrium (random association) with between linked alleles.

A marker in linkage disequilibrium with disease predisposing variants can be particularly useful in detecting susceptibility to disease (or association with other sub-clinical phenotypes) notwithstanding that the marker does not cause the disease. For example, a marker (X) that is not itself a causative element of a disease, but which is in linkage disequilibrium with a gene (including regulatory sequences) (Y) that is a causative element of a phenotype, can be used to indicate susceptibility to the disease in circumstances in which the gene Y may not have been identified or may not be readily detectable. Younger alleles (*i.e.*, those arising from mutation relatively late in evolution) are expected to have a larger genomic segment in linkage disequilibrium. The age of an allele can be determined from whether the allele is shared between different human ethnic human groups and/or between humans and related species.

Unless otherwise apparent from the context, any embodiment, element or feature of the invention can be used in combination with any other.

“Analog” are nucleic acid sequences that have a structure similar to, but not identical to, the native compound but differ from it in respect to certain components. Usually, an analog has the same or similar function to the native compound. Analogs may be synthetic or from a different evolutionary origin and may have a similar or opposite metabolic activity compared to wild type.

Derivatives and analogs may be full length or other than full length. Derivatives or analogs of the nucleic acids of the invention include, nucleic acids that are substantially identical to an exemplified SEQ ID NO. and/or capable of hybridizing to the complement of the sequence under highly stringent, moderately stringent, or low stringent conditions.

The terms "identical" or percent "identity," in the context of two or more nucleic acids or polypeptides, refer to two or more sequences or subsequences that are the same or have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned for maximum correspondence, as measured using a sequence comparison algorithm such as those described below for example, or by visual inspection.

The phrase "substantially identical," in the context of two nucleic acids or polypeptides, refers to two or more sequences or subsequences that have at least 80, preferably at least 85%, more preferably at least 90%, 95%, 99% or higher nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using a sequence comparison algorithm such as those described below for example, or by visual inspection. Preferably, the substantial identity exists over a region of the sequences that is at least 40 residues (*i.e.*, amino acids or nucleotides) in length, preferably over a longer region than 50 residues, more preferably at least about 90-100 residues, and most preferably the sequences are substantially identical over the full length of the sequences being compared, such as the coding region of a nucleotide for example. For example, when a SEQ ID NO. of the invention serves as a reference for comparison with an object nucleic acid, the comparison is preferably performed over the entire length of the SEQ ID NO. of the invention.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (see generally Ausubel *et al.*, *supra*).

Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length *W* in the query sequence, which either match or satisfy some positive-valued threshold score *T* when aligned with a word of the same length in a database sequence. *T* is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters *M* (reward score for a pair of matching residues; always > 0) and *N* (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity *X* from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. For identifying whether a nucleic acid or polypeptide is within the scope of the invention, the default parameters of the BLAST programs are suitable. The BLASTN program (for nucleotide sequences) uses as defaults a word length (*W*) of 11, an expectation (*E*) of 10, *M*=5, *N*=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a word length (*W*) of 3, an expectation (*E*) of 10, and the BLOSUM62 scoring matrix. The TBLASTN program (using protein sequence for nucleotide sequence) uses as defaults a word length (*W*) of 3, an expectation (*E*) of 10, and a BLOSUM 62 scoring matrix. (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)).

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, *e.g.*, Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability ($P(N)$), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under highly stringent conditions. "Bind(s) substantially" refers to complementary hybridization between a probe nucleic acid and a target nucleic acid and embraces minor mismatches that can be accommodated by reducing the stringency of the hybridization media to achieve the desired detection of the target polynucleotide sequence. The phrase "hybridizing specifically to", refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under highly stringent conditions when that sequence is present in a complex mixture (*e.g.*, total cellular) DNA or RNA.

The phrases "specifically binds" refers to a binding reaction which is determinative of the presence of the protein in the presence of a heterogeneous population of proteins and other biologics. Thus, under designated conditions, a specified ligand binds preferentially to a particular protein and does not bind in a significant amount to other proteins present in the sample. A molecule such as antibody that specifically binds to a protein often has an association constant of at least $10^6 M^{-1}$ or $10^7 M^{-1}$, preferably $10^8 M^{-1}$ to $10^9 M^{-1}$, and more preferably, about $10^{10} M^{-1}$ to $10^{11} M^{-1}$ or higher. An antibody that specifically binds to one segment of a protein (*e.g.*, residues 1-10) does not bind to other segments of the protein not included within or overlapping the designated segment.

The term AAA1 gene means nucleotides 163615–684776 of the human genomic contig NT_000380 and allelic or species variants thereof.

The term GPRA gene means nucleotides 471478-691525 of the human genomic contig NT_000380 and allelic or species variants thereof.

A "pharmacological" activity means that an agent exhibits an activity in a screening system that indicates that the agent is or may be useful in the prophylaxis or treatment of a disease. The screening system can be in vitro, cellular, animal or human. Agents can be described as having pharmacological activity notwithstanding that further testing may be required to establish actual prophylactic or therapeutic utility in treatment of a disease.

Pulmonary diseases associated with lower airways obstruction are classified as chronic obstructive pulmonary disease (COPD) or asthma on the basis of the clinical picture.

"Asthma" is a chronic inflammatory disease of the airways causing variable airflow obstruction that is often reversible either spontaneously or with treatment. Asthma is highly correlated with other allergic, IgE mediated diseases, such as allergic rhinitis and dermatitis. COPD is characterized by slowly progressing, mainly irreversible airway obstruction and decreased expiratory flow rate.

DETAILED DESCRIPTION OF THE INVENTION

I. General

This invention is based in part on the discovery and characterization of a novel susceptibility locus for asthma and other IgE mediated diseases, two overlapping genes within the locus, termed GPRA and AAA1, and proteins encoded by the genes. The genes have opposite transcriptional orientations and share intronic regions although not exons. The locus maps within human chromosome 7p15-p14.

Fig. 3 shows the structure of the human GPRA gene. The gene has nine exons, two of which have alternate forms, and eight introns. The total length of the gene from the first base of the first exon to the last base of the last exon is 220047 bp and the coordinates are 471478 (beginning) 691525 (end) in NT_000380. Part of the genomic sequence of GPRA including part of intron 2, exon 3, intron 3, exon 4 and part of intron 4 defines a sublocus referred to as AST1. This locus is 129,017 bp in length. The locus refers to the part of human chromosome 7 having the 129,017 bp of SEQ ID NO:1 or allelic or species variants thereof.

Fig. 13 shows the structure of the human AAA1 gene. The gene has nineteen exons, nine of which have alternate forms, and 18 introns. The total length of the gene from the first base of the first exon to the last base of the last exon is 521,161 bp and the coordinates are 163,615 (beginning) 684776 (end) in NT_000380. Part of the genomic sequence of AAA1 including part of intron 2, exon 3 (a and b), intron 3, exon 4 (a and b), intron 4, exon 5 (a and b), intron 5, exon 6, intron 6, exon 7 (a and b), intron 7, exon 8, intron 8, exon 9, intron 9, exon 10 (a and b) and part of intron 10 occurs within the AST1 locus.

The present application shows that the GPRA and AAA1 genes and AST-1 locus within them are genetically linked and associated with IgE diseases, such as asthma. In addition, the invention provides a collection of polymorphic sites including SNPs (single nucleotide polymorphisms) and DIPs (deletion/insertion polymorphisms) within the locus. Tables 3, 7 and 12 show variant and reference (or wildtype) forms occupying these sites. The variant forms are associated with asthma, other IgE mediated diseases, and/or immune-mediated diseases including autoimmune diseases, and/or cancer and are useful in detection, diagnosis, treatment and prophylaxis thereof.

The AST-1 locus was localized by hierarchical LD mapping in a systemic search for a shared haplotype among the individuals with high IgE serum level across a 19 Mb region that was first identified by a genome wide screen for asthma genes (Laitinen *et al.* 2001). This analysis defined the AST-1 locus between the markers SNP509783 and SNP638799 and provided statistically significant evidence that specific haplotype of 129kb within the linkage peak is a risk factor for asthma related traits (SEQ ID NO:1). The best haplotype patterns of polymorphic sites in GPRA (Table 8) were found with the frequency of 13-18% among the high IgE associated chromosomes compared to that of 3-7% among control chromosomes suggesting the risk ratio of 3.9 [95%CI: 1.8-8.6] for high IgE level (38/304 among affected vs. 7/220 control chromosomes). The best observe associations for polymorphic sites in AAA1 reached the χ^2 values of 8.9-13.6 All markers identified between the positions 509,783-638,799 in the genomic contig NT_000380 were found to be in strong LD and therefore useful for diagnostic purposes.

II. GPRA Polypeptides

The human GPRA gene can be expressed as at least seven different splice variants termed A, B-long, B-short, C, D, E, and F. The cDNAs encoding each form and amino acid sequence of each form are shown in Fig. 4 (SEQ ID NOS:2-15). The different splicing events that lead to the different forms are shown in Fig. 5. It can be seen that the B-long and B-short forms differ from the A form in that the B forms have a different C-terminus encoded by exon 9b, whereas the C terminus of the A form is encoded by exon 9A. The B-short form (SEQ ID NOS:6 and 7) differs from the B-long form in that the B-short form has a deletion of 10 amino acids encoded by exon III. The C-form (SEQ ID NOS:8 and 9) contains a segment encoded by exon 2b which is not found in other forms. The C-form lacks any segments encoded by exons 3-9. The D form (SEQ ID NOS:10 and 11) lacks a segment encoded by exon 3, causing a frameshift, whereby exons 4 and 5 are read in a different reading frame causing truncation within exon 5. Form E (SEQ ID NOS:12 and 13) lacks exon 4 causing a frameshift as a result of which exon 5 is read in a different reading frame causing truncation within exon 5. Form F (SEQ ID NOS:14 and 15) lacks exons 3 and 4.

Fig. 6 shows the expected structural characteristics of the different splice variants. The A form includes 371 amino acids and has an extracellular N-terminal domain, seven transmembrane domains and an intracellular C-terminal domains (SEQ ID NO:3). The transmembrane domains are separated by alternating cytoplasmic and extracellular loops. The B-long form (377 amino acids, SEQ ID NO:5) has a similar structure. The predicted size of the extracellular N-terminal domain is about 50 amino acids anticipating suggesting that the endogenous ligand that interacts with GPRA is a small molecule or peptide. The B-short, C, D, E and F variants of GPRA (SEQ ID NOS: 5, 7, 9, 11, 13, and 15, respectively) lack the 7TM structure present in variants A and B-Long.

Fig. 7 shows the arrangement of the A or B-Long form in a cell membrane. GPRA falls within the conserved class A of G-protein coupled receptors (GPRs) (Rana *et al.* 2001, Johnson *et al.* 2000). GPRs are a large and functionally diverse protein superfamily that form a seven-transmembrane (7TM) helices bundle with alternating extracellular and intracellular loops (see WO 01/18206). Class A contains most well known members of the GPRs such as vasopressin, oxytocin, and bovine rodopsin receptors. For the latter the crystallographic structure is available (Palczewski *et al.* 2000). GPRA is expressed in human tissues consistent with its role in asthma and IgE mediated disease. Transcripts of *GPRA* variants A, B, and C were expressed in several tissues including lung and NCI-H358 cell line that

represent cells of broncho-epithelial origin (Fig. 8 and 9). Based on immunohistochemistry, the A variant of the GPRA protein was expressed in smooth muscle cells of bronchi and arterial wall in human lung as subepithelial of smooth muscle cells in colon (Fig. 10) and the B (-long) variant in the epithelium of bronchi and colon (Fig. 11).

A class of GPRA polypeptides can be defined with reference to the exemplary polypeptides defined by SEQ ID NOS: 3, 5, 7, 9, 11, 13 and 15. The class includes allelic, cognate and induced variants of the exemplified polypeptides. Preferred polypeptides shows substantial sequence identity to one or more of the exemplified polypeptides. The class also includes fragments of the exemplified polypeptides having at least 6, 10, 20, 50 100, 200, or 300 contiguous amino acids from one of the above SEQ ID NOS. Some fragments include one or more isolated domains from one of the exemplified polypeptides, *e.g.*, any of the seven transmembrane domains, any of three extracellular loops, any of three intracellular loops, an N-terminal domain and a C-terminal domain. Some polypeptides comprise the amino acid of one of the above SEQ ID NOS. provided that up to 1, 2, 5, 10, 20, 30 or 34 amino acids can be inserted, deleted or substituted relative to the SEQ ID NO. Some GPRA polypeptides comprise an epitope found in one of the exemplified SEQ ID NOS. but not in others. For example, B-Long (SEQ ID NO:5) has a C-terminal epitope of 35 amino acids not found in any of the other forms. Some GPRA polypeptides contain a polymorphic site shown in Table 7 occupied by a variant form.

Preferred GPRA polypeptides are receptors that can be activated to transduce a signal by the same ligand as the exemplified GPRA polypeptide of SEQ ID NOS: 3, 5, 7, 9, 11, 13, and 15. Binding of a ligand or analog to a G-protein receptor induces an alteration in receptor G-protein interaction. The receptor G-protein interaction releases GDP specifically bound to the G protein and permits the binding of GTP, which activates the G protein. Activated G-protein dissociates from the receptor and activates an effector protein, which in turn regulates intracellular levels of second messengers, such as adenylyl cyclase, guanylyl cyclase, and phospholipase C. Signal transduction can be monitored in cells transfected with a reporter construct consisting of several copies of the consensus cAMP response element (Arias *et al.*, *Nature* 370, 226-229 (1994)), a minimal tk promoter, and a secreted alkaline phosphatase gene (Clontech) or can be monitored in changes of the concentrations of intracellular calcium.

III. AAA1 Polypeptides

The human AAA1 gene can be expressed as at least thirteen different splice variants termed IA, IB, II, III, IVA, IVB, V, VI, VII, VIII, IX, X, and XI. The cDNAs encoding each form and amino acid sequence of each form are shown in Fig. 20 (SEQ ID NOS:16-41). The different splicing events that lead to the different forms are shown in Fig. 14. It can be seen that all splice variants have exon VI and the same core peptide sequence (AYVRRNAGRQFSHCNLHAHQFLVRRKQ). The AAA1 gene is expressed predominantly in the testis, brain, placenta, lung, heart, skeletal muscle, kidney, liver, fetal liver and fetal lung. In addition to the variants described in Fig. 14, using 3'-RACE we have found also several other variants expressed in different tissues. They all share the exon six.

A class of AAA1 polypeptides can be defined with reference to the exemplary polypeptides defined by SEQ ID NOS:17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41. The class includes allelic, cognate and induced variants of the exemplified polypeptides. Preferred polypeptides shows substantial sequence identity to one or more of the exemplified polypeptides. The class also includes fragments of the exemplified polypeptides having at least 6, 10, 20, or 50 contiguous amino acids from one of the above SEQ ID NOS. Some fragments include or consist of the above core peptide sequence. Some polypeptides comprise the amino acid of one of the above SEQ ID NOS. provided that up to 1, 2, 5, 10, or 20 amino acids can be inserted, deleted or substituted relative to the SEQ ID NO. Some AAA1 polypeptides comprise an epitope found in one of the exemplified SEQ ID NOS. but not in others. For example, only the variant 1B contains the peptide sequence encoded by exon III.

IV. GPRA Nucleic Acids

The present invention provides isolated nucleic acids encoding GPRA polypeptides. The nucleic acids can be, for example, genomic, cDNA, RNA or mini-gene (*i.e.*, a hybrid of genomic and cDNA). Genomic and mini-gene nucleic acids contain at least one intronic segment from a GPRA gene, meaning the section of human chromosome 7p15-p14 having the structure shown in Fig. 3, or allelic and cognate variants thereof. Nucleic acids of the invention can include coding regions, intronic regions, 3' untranslated regions, 5' untranslated regions, 3' flanking regions, 5' flanking regions, enhancers, promoters and other regulatory sequences. A class of GPRA nucleic acids can be defined with reference to exemplified GPRA polypeptide and nucleic acid sequences. The class includes allelic, cognate and

induced variants of the exemplified sequences, as well as nucleotide substitutions, which due to the degeneracy of the genetic code, do not effect the amino acid sequence of an encoded polypeptide. The class includes nucleic acids that encode the GPRA polypeptides described above. The class also includes GPRA nucleic acids showing substantial sequence identity to exemplified GPRA nucleic acids (*i.e.*, SEQ ID NOS:1, 2, 4, 6, 8, 10, 12 and 14). SEQ ID NO:1 is a genomic sequence representing the part of a GPRA gene designated Ast-1 and shown in Fig. 3. SEQ ID NO:2 is a 1113 bp cDNA encoding GPRA variant A (Fig. 4A). SEQ ID NO:4 is a 1132 bp cDNA encoding GPRA variant B-Long (Fig. 4B). SEQ ID NOS:6, 8, 10, 12, and 14 encode GPRA variants B-Short, C, D, E and F.

The class also includes GPRA nucleic acids that hybridize under highly stringent conditions with at least one of the exemplified nucleic acids. Some GPRA nucleic acids of the invention hybridize under highly stringent conditions with at least one of the GPRA nucleic acids of the invention without hybridizing under the same conditions to others. For example, some GPRA nucleic acids hybridize to at least one of SEQ ID NOS:1, 4, 6, 8, 10, 12 and 14 without hybridizing to SEQ ID NO:2. Some GPRA nucleic acids include a polymorphic site occupied by a variant form as shown in Table 3 or Table 7. Some GPRA nucleic acids are genomic or minigene. Such nucleic acids contain at least one intronic sequence from any of the intron of a GPRA gene. Inclusion of an intronic sequence can be useful for increasing expression levels of a nucleic acid in cells or transgenic animals. Inclusion of an intronic sequence from introns 2, 3 or 4 is particularly useful for analyzing splice variation of the GPRA gene. Nucleic acids also include fragments of exemplified sequences SEQ ID NOS:1, 4, 6, 8, 10, 12 and 14. The fragments typically contain up to 10, 20, 25, 50, 100, 300, 400, 500 or 1000 or 1500 nucleic acids from any of the above SEQ ID NOS. Optionally, fragments include a polymorphic site shown in Table 3 or 7. The polymorphic site can be occupied by either a reference or variant form.

GPRA nucleic acids can be linked to other nucleic acids with which they are not naturally associated such as vector or a heterologous promoter sequence. Preferred nucleic acids of the invention include or are immediately adjacent to at least one of the polymorphic sites shown in Table 3 or 7. Such nucleic acids are useful as primers or probes for detection of specific alleles. Other nucleic acids are useful for expressing proteins encoded by the AST-1 locus in cells or transgenic animals. Other nucleic acids are useful for achieving gene suppression either by homologous recombination to generate a knockout animal or an antisense or siRNA

mechanism. Nucleic acids are also useful for identifying, purifying, and isolating nucleic acids encoding other, non-human, mammalian forms of asthma locus-1.

Nucleic acids of the invention can be isolated using standard molecular biology techniques and the provided sequence information (Ausubel *et al*, In Current protocols in Molecular Biology, John Wiley and Sons, publishers, 1989); Sambrook *et al*, supra). Cognate forms (*i.e.*, nucleic acids encoding asthma locus-1 molecules derived from species other than human) or other related sequences (*e.g.*, paralogs) can be obtained by low, moderate or high stringency hybridization with all or a portion of the particular human sequence as a probe using methods well known in the art for nucleic acid hybridization and cloning.

V. AAA1 Nucleic Acids

The present invention provides isolated nucleic acids encoding AAA1 polypeptides. The nucleic acids can be, for example, genomic, cDNA, RNA or mini-gene (*i.e.*, a hybrid of genomic and cDNA). Genomic and mini-gene nucleic acids contain at least one intronic segment from an AAA1 gene, meaning the section of human chromosome 7p15-p14 having the structure shown in Fig. 13, or allelic and cognate variants thereof. Nucleic acids of the invention can include coding regions, intronic regions, 3' untranslated regions, 5' untranslated regions, 3' flanking regions, 5' flanking regions, enhancers, promoters and other regulatory sequences. A class of AAA1 nucleic acids can be defined with reference to exemplified AAA1 polypeptide and nucleic acid sequences. The class includes allelic, cognate and induced variants of the exemplified sequences, as well as nucleotide substitutions, which due to the degeneracy of the genetic code, do not effect the amino acid sequence of an encoded polypeptide. The class includes nucleic acids that encode the AAA1 polypeptides described above. The class also includes AAA1 nucleic acids showing substantial sequence identity to exemplified AAA1 nucleic acids (*i.e.*, SEQ ID NOS:1, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40). SEQ ID NO:1 is a genomic sequence representing the part of an AAA1 gene designated Ast-1 and shown in Fig. 13. SEQ ID NOS:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40 encode cDNA splice variants shown in Fig. 20.

The class also includes AAA1 nucleic acids that hybridize under highly stringent conditions with at least one of the exemplified nucleic acids. Some AAA1 nucleic acids of the invention hybridize under highly stringent conditions with at least one of the exemplified nucleic acids (*i.e.*, SEQ ID NOS:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40) without hybridizing

under the same conditions to other exemplified nucleic acids. Some AAA1 nucleic acids include a polymorphic site occupied by a variant form as shown in Table 12. Some AAA1 nucleic acids are genomic or minigene. Such nucleic acids contain at least one intronic sequence from any of the introns of an AAA1 gene. Nucleic acids also include fragments of exemplified sequences SEQ ID NOS:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40. The fragments typically contain up to 10, 20, 25, 50, or 100, nucleotides from any of the above SEQ ID NOS. Optionally, fragments include a polymorphic site shown in Table 12. The polymorphic site can be occupied by either a reference or variant form.

AAA1 nucleic acids can be linked to other nucleic acids with which they are not naturally associated such as vector or a heterologous promoter sequence. Preferred nucleic acids of the invention include or are immediately adjacent to at least one of the polymorphic sites shown in Table 3 or Table 12. Such nucleic acids are useful as primers or probes for detection of specific alleles. Other nucleic acids are useful for expressing AAA1 polypeptides in cells or transgenic animals. Other nucleic acids are useful for achieving gene suppression either by homologous recombination to generate a knockout animal or an antisense or siRNA mechanism. Nucleic acids are also useful for identifying, purifying, and isolating nucleic acids encoding other, non-human, mammalian forms of the AAA1 gene.

As well as or instead of encoding proteins, AAA1 transcripts can have roles in regulating expression of GPRA nucleic acids. A class of noncoding (ncRNA) has been described with regulatory activity in gene transcription and splicing (see e.g. reviews by Michel 2002; Numata *et al.* 2003).

VI. Antibodies

The GPRA and AAA1 polypeptides of the invention are useful for generating antibodies. The antibodies can be polyclonal antibodies, distinct monoclonal antibodies or pooled monoclonal antibodies with different epitopic specificities. Monoclonal antibodies are made from antigen-containing fragments of the protein by standard procedures according to the type of antibody (see, e.g., Kohler, *et al.*, *Nature*, 256:495, (1975); and Harlow & Lane, *Antibodies, A Laboratory Manual* (C.S.H.P., NY, 1988) Queen *et al.*, *Proc. Natl. Acad. Sci. USA* 86:10029-10033 (1989) and WO 90/07861; Dower *et al.*, WO 91/17271 and McCafferty *et al.*, WO 92/01047 (each of which is incorporated by reference for all purposes). Phage display technology can also be used to mutagenize CDR regions of antibodies previously

shown to have affinity for the peptides of the present invention. Some antibodies bind to an epitope present in one form of GPRA or AAA1 but not others. For example, some antibodies bind to an epitope within amino acids 343-377 of GPRA long B (SEQ ID NO:4). Some antibodies specifically bind to a GPRA polypeptide containing a variant form at a polymorphic site shown in Table 7 without binding to polypeptides containing a reference form at the site. The antibodies can be purified, for example, by binding to and elution from a support to which the polypeptide or a peptide to which the antibodies were raised is bound.

Antibodies of the invention are useful, for example, in screening cDNA expression libraries and for identifying clones containing cDNA inserts which encode structurally-related, immunocrossreactive proteins. See, for example, Aruffo & Seed, *Proc. Natl. Acad. Sci. USA* 84:8573-8577 (1977) (incorporated herein by reference in its entirety for all purposes).

Antibodies are also useful to identify and/or purify immunocrossreactive proteins that are structurally related to SEQ ID NOS:3, 5, 7, 9, 11, 13 and 15 and 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 and fragments thereof used to generate the antibody.

VII. Polymorphisms of the Invention

The present invention provides a collection of polymorphic sites both single nucleotide polymorphisms (SNPs) and deletion/insertion polymorphisms (DIPs) in the GPRA gene. SNPs and DIPs can be used in mapping the human genome and, when a SNP or a DIP is linked with a disease or condition, to clarify genetic basis of the disease or condition, in this particular case, at least of asthma.

Table 3 describes 169 SNPs and 18 DIPs occurring within exons 3 and 4 and introns 2, 3 and 4 of a GPRA gene having variant forms associated with asthma. For AAA1 the corresponding polymorphisms are located in exons 4, 6, 9, 10a /b and introns 2, 3, 4, 5, 6, 7, 8, 9, and 10 (Figure 13). The locus containing these polymorphisms is also referred to as AST-1 (SEQ ID NO:1). Some of the variant forms of these polymorphism may associate with asthma due to an effect on the splicing of GPRA and/or AAA1 gene in the formation of a transcript. Other polymorphic sites may be in linkage disequilibrium with such sites or sites in coding regions that exert effects through changing the activity of GPRA or AAA1 polypeptides. Seven different haplotypes H1-H7 were identified that explain 94.5% of the genetic variation detected at the AST-1 locus and are described in Table 3. Haplotype H1 was found identical to the reference sequence NT_000380. The first column in Table 3 indicates

the types of polymorphism (*e.g.*, SNP or insertion/deletion). The second column indicates the position that the polymorphism occurs in SEQ ID NO:1. The third column indicates the position of the polymorphism occurs in NT_000380. The fourth column indicates several contiguous nucleotides flanking the polymorphic sites and the nature of the variant (*i.e.*, asthma associated) and reference/wildtype polymorphic forms. In all polymorphic sites, the allele present in SEQ ID NO:1 and NT_000380 is indicated first, and the allele from another source second. Columns 5-11 indicate alleles present at that site in haplotypes H1-H7. Column 12 indicates the location of polymorphisms in GPRA and AAA1 exons. In the AST-1 locus, there are coding polymorphism in AAA1 gene (exon 6, Asn ->Ser) and one in GPRA gene (exon 3, Asn -> Ile). Table 13 shows polymorphisms that are unique for haplotypes H1-H7, respectively, and Table 14 polymorphisms that can be employed to recognize specific haplotype combinations.

The SNPs include the following substitutions with the variant form being indicated before the reference form of SEQ ID NO: 1. Position 1 (preferably T (*i.e.*, variant form) instead of C (reference form)), position 93 (preferably G instead of A), position 918 (preferably A instead of G), position 983 (preferably T instead of A), position 987 (preferably C instead of T), position 1542 (preferably T instead of C), position 1710 (preferably G instead of A), position 1818 (preferably T instead of C), position 1927 (preferably T instead of A), position 2254 (preferably C instead of T), position 2937 (preferably A instead of G), position 3877 (preferably C instead of G), position 4012 (preferably A instead of C), position 4631 (preferably C instead of T), position 4689 (preferably G instead of C), position 4961 (preferably G instead of A), position 5442 (preferably G instead of C), position 5634 (preferably A instead of G), position 5850 (preferably G instead of A), position 6312 (preferably T instead of C), position 6392 (preferably T instead of G), position 6485 (preferably A instead of G), position 6522 (preferably G instead of C), position 6646 (preferably G instead of A), position 6739 (preferably A instead of G), position 6760 (preferably C instead of T), position 7125 (preferably A instead of C), position 7229 (preferably T instead of C), position 7277 (preferably G instead of C), position 7303 (preferably T instead of G), position 7305 (preferably C instead of G), position 7496 (preferably T instead of C), position 7550 (preferably A instead of G), position 8490 (preferably T instead of C), position 9649 (preferably G instead of T), position 10816 (preferably C instead of T), position 11858 (preferably G instead of A), position 12581 (preferably C instead of G), position 16845 (preferably C instead of T), position 16893

(preferably C instead of T), position 16980 (preferably C instead of T), position 17147
(preferably A instead of G), position 17209 (preferably A instead of C), position 17435
(preferably G instead of A), position 18383 (preferably A instead of G), position 18927
(preferably T instead of G), position 18978 (preferably G instead of A), position 19268
(preferably C instead of G), position 19272 (preferably T instead of A), position 19360
(preferably A instead of G), position 19452 (preferably A instead of G), position 19671
(preferably A instead of G), position 19712 (preferably A instead of C), position 19774
(preferably A instead of C), position 20038 (preferably C instead of T), position 20089
(preferably A instead of T), position 20309 (preferably A instead of G), position 20395
(preferably C instead of T), position 20789 (preferably T instead of G), position 21850
(preferably T instead of C), position 22475 (preferably C instead of T), position 22493
(preferably G instead of A), position 22715 (preferably A instead of G), position 22869
(preferably C instead of T), position 22934 (preferably T instead of A), position 24007
(preferably C instead of T), position 24264 (preferably T instead of G), position 24869
(preferably C instead of T), position 26198 (preferably T instead of C), position 26356
(preferably T instead of C), position 26675 (preferably G instead of A), position 27404
(preferably T instead of C), position 28197 (preferably A instead of G), position 28770
(preferably T instead of C), position 28785 (preferably A instead of G), position 28858
(preferably C instead of T), position 28866 (preferably C instead of G), position 31224
(preferably A instead of G), position 31910 (preferably A instead of G), position 32124
(preferably A instead of C), position 32185 (preferably C instead of T), position 32976
(preferably C instead of T), position 33350 (preferably A instead of C), position 33798
(preferably A instead of G), position 34362 (preferably C instead of G), position 34716
(preferably C instead of A), position 35559 (preferably T instead of C), position 36551
(preferably A instead of G), position 36909 (preferably A instead of C), position 37327
(preferably T instead of G), position 37415 (preferably A instead of G), position 37685
(preferably G instead of A), position 37931 (preferably T instead of C), position 37959
(preferably T instead of C), position 39314 (preferably A instead of G), position 39343
(preferably T instead of G), position 39927 (preferably C instead of T), position 45826
(preferably T instead of C), position 50197 (preferably T instead of G), position 50334
(preferably A instead of G), position 50493 (preferably A instead of G), position 50632
(preferably A instead of G), position 50835 (preferably C instead of A), position 50955
(preferably C instead of G), position 51217 (preferably A instead of G), position 51476
(preferably G instead of A), position 51536 (preferably T instead of C), position 51861

(preferably G instead of C), position 51884 (preferably T instead of G), position 51975 (preferably C instead of G), position 52573 (preferably A instead of G), position 52776 (preferably C instead of G), position 53803 (preferably T instead of C), position 53922 (preferably C instead of T), position 54148 (preferably A instead of G), position 54199 (preferably C instead of T), position 54641 (preferably C instead of G), position 54751 (preferably C instead of T), position 55000 (preferably A instead of G), position 55134 (preferably A instead of G), position 56683 (preferably T instead of C), position 56856 (preferably T instead of C), position 57790 (preferably C instead of A), position 60559 (preferably G instead of C), position 60604 (preferably T instead of A), position 61165 (preferably A instead of G), position 64559 (preferably T instead of C), position 65171 (preferably G instead of A), position 65857 (preferably G instead of T), position 66164 (preferably T instead of C), position 66190 (preferably T instead of C), position 66526 (preferably T instead of C), position 66902 (preferably G instead of A), position 67857 (preferably A instead of T), position 67919 (preferably C instead of T), position 72270 (preferably A instead of G), position 75115 (preferably A instead of G), position 76080 (preferably T instead of C), position 76101 (preferably C instead of G), position 81912 (preferably T instead of A), position 82203 (preferably G instead of A), position 82332 (preferably T instead of C), position 82922 (preferably A instead of T), position 83552 (preferably T instead of C), position 85227 (preferably C instead of T), position 85271 (preferably A instead of G), position 107610 (preferably C instead of T), position 110989 (preferably C instead of T), position 111012 (preferably T instead of C), position 112030 (preferably C instead of G), position 112037 (preferably G instead of T), position 112283 (preferably T instead of A), position 112726 (preferably A instead of T), position 112859 (preferably C instead of A), position 113428 (preferably G instead of C), position 113645 (preferably T instead of A), position 113944 (preferably G instead of T), position 114945 (preferably G instead of A), position 115192 (preferably G instead of C), position 115628 (preferably C instead of T), position 116032 (preferably A instead of G), position 116464 (preferably A instead of G), position 116515 (preferably A instead of G), position 116926 (preferably C instead of T), position 117276 (preferably A instead of T), position 123667 (preferably A instead of G), position (preferably 123770 instead G of A), position 123788 (preferably C instead of G), and position 129017 (preferably G instead of A).

DIPS of the invention defined relative to SEQ ID NO: 1 include: positions 184-185

(preferably insertion of AAGATA), positions 1202-1206 (preferably deletion of TAAGT),

positions 6786-6817 (preferably repeat of (TAAA)₇), position 6821 (preferably deletion of T), positions 7240-7243 (preferably deletion of ACTT), positions 7306-7308 (preferably deletion of TGT), positions 7334-7335 (preferably deletion of AT), positions 9012-9035 (preferably repeat of (CT)₁₀), positions 9199-9201 (preferably deletion of TCT), : position 9355-9356 (preferably insertion of T), positions 9782-9785 (preferably deletion of GTCT), positions 22122-22123 (preferably deletion of TG), positions 26929-26968 (preferably repeat of (TAAA)₁₁), positions 28495-28564 (preferably repeat of (CA)₁₂), position 34909 (preferably deletion of T), positions 38850-38852 (preferably deletion of CTC), positions 51022-51049 (preferably repeat of (CA)₈), and positions 52286-52287 (preferably insertion of CC).

Table 7 provide 13 additional polymorphisms occurring within exonic regions of a GPRA gene. The next eight columns indicate the position of a polymorphic site in SEQ ID NO:1, 2, 4, 6, 8, 10, 12 or 14 respectively. The next column indicates a polymorphic site and flanking nucleotides. The form of the polymorphic site present in one or more of SEQ ID NOS:1, 2, 4, 6, 8, 10, 12, and 14 is shown first and the form present in another source is indicated second. In all cases except those indicated by a *, the polymorphic form indicated first is the wildtype or reference form and the polymorphic form indicated second is the asthma or IgE associated form. In the three polymorphic sites indicated with a *, the variant form of the polymorphic site is found in NT_000380 (and SEQ ID NO:1, 2, 4, 6, 8, 10, 12 and/or 14), and the reference or wildtype allele is found in a different individual. The last column of the table shows whether an amino acid change occurs due to the polymorphisms. The positions of these polymorphisms defined with respect to SEQ ID NO:2 are: position 448 (T (variant) preferably to A (wildtype or reference)), position 776 (C preferably to T), position 851 (C preferably to G), position 1159 (G preferably to A), position 1199 (T preferably to C), and position 1529 (C preferably to T); or in any one of the following positions as defined by SEQ ID NO:4: position 1206 (A preferably to C), position 1225 (T preferably to C), position 1330 (T preferably to A), and position 1338 (G preferably to A); or in any one of the following positions as defined by SEQ ID NO:8: position 585 (T preferably to C), position 655 (C preferably to A), and position 681 (A preferably to G).

Table 8 shows the five polymorphic sites within Table 7 that showed the highest association with high serum IgE level. Two of the polymorphic sites involved nonconservative amino acid changes, one within the important region of exon 3 within AST-1. Two of the

polymorphic sites were within coding regions but did not cause amino acid changes. These sites may show strong association as a result of linkage disequilibrium with causative polymorphic sites. A fifth polymorphic site is within the 3' UTR of exon 9A. This site may show a strong association for the same reason or because the nucleotide substitution has an effect on mRNA stability.

Table 12 shows associations for five polymorphic sites occurring within the AAA1 gene. At each site, the wildtype polymorphic form is indicated first, and the variant form second. The table shows relative frequency of associating haplotypes in individuals with high serum IgE level compared to those in control individuals with low serum IgE level. All haplotypes shown in the table gave chi square values of at least 8.9 ($P < 0.01$) for association.

VIII. Analysis of Polymorphisms

A. Preparation of Samples

Polymorphisms are detected in a target nucleic acid from an individual being analyzed. For assay of genomic DNA, virtually any biological sample (other than pure red blood cells) is suitable. For example, convenient tissue samples include whole blood, semen, saliva, tears, urine, fecal material, sweat, buccal, skin and hair. For assay of cDNA or mRNA, the tissue sample must be obtained from an organ in which the target nucleic acid is expressed.

Many of the methods described below require amplification of DNA from target samples. This can be accomplished by *e.g.*, PCR. See generally PCR Technology: Principles and Applications for DNA Amplification (ed. H.A. Erlich, Freeman Press, NY, NY, 1992); PCR Protocols: A Guide to Methods and Applications (eds. Innis, *et al.*, Academic Press, San Diego, CA, 1990); Mattila *et al.*, Nucleic Acids Res. 19, 4967 (1991); Eckert *et al.*, PCR Methods and Applications 1, 17 (1991); PCR (eds. McPherson *et al.*, IRL Press, Oxford); and U.S. Patent 4,683,202 (each of which is incorporated by reference for all purposes).

Other suitable amplification methods include the ligase chain reaction (LCR) (see Wu and Wallace, Genomics 4, 560 (1989), Landegren *et al.*, Science 241, 1077 (1988), transcription amplification (Kwoh *et al.*, Proc. Natl. Acad. Sci. USA 86, 1173 (1989)), and self-sustained sequence replication (Guatelli *et al.*, Proc. Nat. Acad. Sci. USA, 87, 1874 (1990)) and nucleic acid based sequence amplification (NASBA). The latter two amplification methods involve isothermal reactions based on isothermal transcription, which produce both single stranded

RNA (ssRNA) and double stranded DNA (dsDNA) as the amplification products in a ratio of about 30 or 100 to 1, respectively.

B. Detection of Polymorphisms in Target DNA

The identity of bases occupying the polymorphic sites shown in Table 3, 7 or 12 can be determined in an individual (*e.g.*, a patient being analyzed) by several methods, which are described in turn.

1. Single Base Extension Methods

Single base extension methods are described by *e.g.*, US 5,846,710, US 6,004,744, US 5,888,819 and US 5,856,092. In brief, the methods work by hybridizing a primer that is complementary to a target sequence such that the 3' end of the primer is immediately adjacent to but does not span a site of potential variation in the target sequence. That is, the primer comprises a subsequence from the complement of a target polynucleotide terminating at the base that is immediately adjacent and 5' to the polymorphic site. The hybridization is performed in the presence of one or more labeled nucleotides complementary to base(s) that may occupy the site of potential variation. For example, for a biallelic polymorphisms two differentially labeled nucleotides can be used. For a tetraallelic polymorphism four differentially labeled nucleotides can be used. In some methods, particularly methods employing multiple differentially labeled nucleotides, the nucleotides are dideoxynucleotides. Hybridization is performed under conditions permitting primer extension if a nucleotide complementation a base occupying the site of variation in the target sequence is present. Extension incorporates a labeled nucleotide thereby generating a labeled extended primer. If multiple differentially labeled nucleotides are used and the target is heterozygous then multiple differentially labeled extended primers can be obtained. Extended primers are detected providing an indication of which base(s) occupy the site of variation in the target polynucleotide. The methods are particularly useful for SNPs.

2. Allele-Specific Probes

The design and use of allele-specific probes for analyzing polymorphisms is described by *e.g.*, Saiki *et al.*, Nature 324, 163-166 (1986); Dattagupta, EP 235,726, Saiki, WO 89/11548. Allele-specific probes can be designed that hybridize to a segment of target DNA from one individual but do not hybridize to the corresponding segment (corresponding segments being defined by maximum alignment of the sequences being compared) from another individual

due to the presence of different polymorphic forms in the respective segments from the two individuals. Hybridization conditions should be sufficiently stringent that there is a significant difference in hybridization intensity between alleles, and preferably an essentially binary response, whereby a probe hybridizes to only one of the alleles. Some probes are designed to hybridize to a segment of target DNA such that the polymorphic site aligns with a central position (*e.g.*, in a 15 mer at the 7 position; in a 16 mer, at either the 8 or 9 position) of the probe. This design of probe achieves good discrimination in hybridization between different allelic forms.

Allele-specific probes are often used in pairs, one member of a pair showing a perfect match to a reference form of a target sequence and the other member showing a perfect match to a variant form. Several pairs of probes can then be immobilized on the same support for simultaneous analysis of multiple polymorphisms within the same target sequence.

3. Tiling Arrays

SNPs and insertion deletion polymorphism can be detected by hybridizing target nucleic acids to arrays of probes tiling a reference sequence or one or more variant forms of the reference sequence as described by WO 95/11995 (incorporated by reference in its entirety for all purposes).

4. Allele-Specific Amplification Methods

An allele-specific primer hybridizes to a site on target DNA overlapping a polymorphism and only primes amplification of an allelic form to which the primer exhibits perfect complementarity. See Gibbs, *Nucleic Acid Res.* 17, 2427-2448 (1989). This primer is used in conjunction with a second primer which hybridizes at a distal site. Amplification proceeds from the two primers leading to a detectable product signifying the particular allelic form is present. A control is usually performed with a second pair of primers, one of which shows a single base mismatch at the polymorphic site and the other of which exhibits perfect complementarity to a distal site. The single-base mismatch prevents amplification and no detectable product is formed. In some methods, the mismatch is included in the 3'-most position of the oligonucleotide aligned with the polymorphism because this position is most destabilizing to elongation from the primer. See, *e.g.*, WO 93/22456.

5. Direct-Sequencing

The direct analysis of the sequence of polymorphisms of the present invention can be accomplished using either the dideoxy- chain termination method or the Maxam -Gilbert method (see Sambrook *et al.*, Molecular Cloning, A Laboratory Manual (2nd Ed., CSHP, New York 1989); Zyskind *et al.*, Recombinant DNA Laboratory Manual, (Acad. Press, 1988)).

6. Denaturing Gradient Gel Electrophoresis

Amplification products generated using the polymerase chain reaction can be analyzed by the use of denaturing gradient gel electrophoresis. Different alleles can be identified based on the different sequence-dependent melting properties and electrophoretic migration of DNA in solution. Erlich, ed., PCR Technology, Principles and Applications for DNA Amplification, (W.H. Freeman and Co, New York, 1992), Chapter 7.

7. Single-Strand Conformation Polymorphism Analysis

Alleles of target sequences can be differentiated using single-strand conformation polymorphism analysis, which identifies base differences by alteration in electrophoretic migration of single stranded PCR products, as described in Orita *et al.*, Proc. Nat. Acad. Sci. 86, 2766-2770 (1989). Amplified PCR products can be generated as described above, and heated or otherwise denatured, to form single stranded amplification products. Single-stranded nucleic acids may refold or form secondary structures which are partially dependent on the base sequence. The different electrophoretic mobilities of single-stranded amplification products can be related to base-sequence difference between alleles of target sequences.

8. Other Methods

Include oligonucleotide ligation assay or restriction fragment length polymorphism (RFLP) (see, for example, Current Protocols in Molecular Biology, eds. Ausubel *et al.*, John Wiley & Sons:1992, and Landegren *et al.*, "Reading Bits of Genetic Information: Methods for Single-Nucleotide Polymorphism Analysis", Genome Research 8:769-776).

VII. Methods of Use

A. Discovery of New Polymorphic Sites in AST-1 Locus

Additional polymorphic sites beyond those listed in Tables 3, 7 and 12 within GPRA and AAA1 genes and particularly, the AST-1 locus thereof can be identified by comparing a

GPRA or AAA1 nucleic acid from different individuals with a reference sequence, such as SEQ. ID NO:1, 2, 4, 6, 8, 10, 12 or 14 or SEQ ID NOS:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 or 40. Preferably, the different individuals suffer or at risk of an immune disorder, such as autoimmune disease, an IgE-mediated disorder, such as asthma, chronic obstructive pulmonary disease or cancer. If a new polymorphic site (*e.g.*, SNP) is found, then the link with a specific disease can be determined by comparing the frequency of variant polymorphic forms occupying this site in populations of patients with or without a disease being analyzed (or susceptibility thereto). A polymorphic site or region may be located in any part of the locus, *e.g.*, exons, introns and promoter regions of genes.

B. Diagnostic and Detection Methods

The present invention further provides means for prognostic or diagnostic assays for determining if a subject has or is increasingly likely to develop a disease associated with the variation or dysfunction of the asthma locus of the invention. Such diseases include in addition to asthma, other IgE mediated diseases, autoimmune diseases, atopic and other immune diseases, chronic obstructive pulmonary disease and cancer. Basically, such assays comprise a detection step, wherein the presence or absence of a variant polymorphic form in the asthma locus-1 relative to SEQ ID NO:1, 2, 4, 6, 8, 10, 12, or 14 or SEQ ID NO: 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40 is determined in a biological sample from the subject. The detection step can be performed by any of the methods described above. Analogous assays can be performed in which variations in protein sequence relative to SEQ ID NOS:3, 5, 7, 9, 11, 13 or 15 or SEQ ID NO: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 are determined.

In particular, the present invention is directed to a method of determining the presence or absence of a variant form of a polymorphic site of the invention in a biological sample from a human for diagnosis of asthma or for assessing the predisposition of an individual to asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer. The method comprises determining the sequence of the nucleic acid of a human at one or more the above SNP or DIP positions shown in Tables 3, 7 and 12. Optionally, the sample is contacted with oligonucleotide primers so that the nucleic acid region containing the potential single nucleotide polymorphism is amplified by polymerase chain reaction prior to determining the sequence.

The polymorphic sites can be analyzed individually or in sets for diagnostic and prognostic purposes. The conclusion drawn from the analysis depends on the nature and number of polymorphic sites analyzed. Some polymorphic sites have variant polymorphic forms that are causative of disease. Detection of such a polymorphic form provides at least a strong indication of presence or susceptibility to disease. Other polymorphic sites have variant polymorphic forms that are not causative of disease but are in equilibrium dislinkage with a polymorphic form that is causative. Detection of noncausative polymorphic forms provides an indication of risk of presence or susceptibility to disease. Detection of multiple variant forms at several polymorphic sites in a GPRA and or AAA1 gene and particularly the parts of these genes within the AST-1 locus thereof provides an indication of increased risk of presence or susceptibility to disease. The results from analyzing the polymorphic sites of the invention can be combined with analysis of other loci that associate with the same disease (e.g., asthma) as polymorphic sites of the invention. Alternatively, or additionally risk of disease can be confirmed by performing conventional medical diagnostic tests of patient symptoms.

C. Clinical Trials

The polymorphisms of the invention are also useful for conducting clinical trials of drug candidates for the diseases noted above particularly asthma. Such trials are performed on treated or control populations having similar or identical polymorphic profiles at a defined collection of polymorphic sites. Use of genetically matched populations eliminates or reduces variation in treatment outcome due to genetic factors, leading to a more accurate assessment of the efficacy of a potential drug.

Furthermore, the polymorphisms of the invention may be used after the completion of a clinical trial to elucidated differences in response to a given treatment. For example, the set of polymorphisms may be used to stratify the enrolled patients into disease sub-types or classes. It may further be possible to use the polymorphisms to identify subsets of patients with similar polymorphic profiles who have unusual (high or low) response to treatment or who do not respond at all (non-responders). In this way, information about the underlying genetic factors influencing response to treatment can be used in many aspects of the development of treatment (these range from the identification of new targets, through the design of new trials to product labeling and patient targeting). Additionally, the polymorphisms may be used to identify the genetic factors involved in adverse response to treatment (adverse events). For

example, patients who show adverse response may have more similar polymorphic profiles than would be expected by chance. This allows the early identification and exclusion of such individuals from treatment. It also provides information that can be used to understand the biological causes of adverse events and to modify the treatment to avoid such outcomes.

VIII. Kits

The invention also features diagnostic or prognostic kits for use in detecting the presence of a SNP in a GPRA and/or AAA1 gene or particularly an AST-1 locus thereof in a biological sample. The kit provides means for the diagnostics of asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer or for assessing the predisposition of an individual to a disease mediated by variation or dysfunction of a GPRA and/or AAA1 gene. The kit can comprise a labeled compound capable of detecting the nucleic acid of a GPRA and/or AAA1 gene in a biological sample. The kit can also comprise nucleic acid primers or probes capable of hybridizing specifically to at least a portion of a GPRA gene. Preferably, the primer is a minisequencing primer specific to any one of the above polymorphisms. The kit can be packaged in a suitable container and preferably it contains instructions for using the kit.

Often, the kits contain one or more pairs of allele-specific oligonucleotides hybridizing to different forms of a polymorphism. In some kits, the allele-specific oligonucleotides are provided immobilized to a substrate. For example, the same substrate can comprise allele-specific oligonucleotide probes for detecting at least 10, 50 or all of the polymorphisms shown in Tables 3, 7, 12, 13, and 14. Optional additional components of the kit include, for example, restriction enzymes, reverse-transcriptase or polymerase, the substrate nucleoside triphosphates, means used to label (for example, an avidin-enzyme conjugate and enzyme substrate and chromogen if the label is biotin), and the appropriate buffers for reverse transcription, PCR, or hybridization reactions.

IX. Transgenic Animals and Cells

The present invention also makes available plasmids and vectors comprising the nucleic acid of the invention, which constructs can be used when studying biological activities of the AST-1 locus. The selecting of a suitable plasmid or vector for a certain use is within the abilities of a skilled artisan. As the host cell may be any prokaryotic or eukaryotic cell, a plasmid or vector comprising the locus can be used in the duplication of the locus (see, for

example, Sambrook *et al*, "Molecular Cloning: A Laboratory Manual", Second Edition, Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (1989)).

Nucleic acids of the invention can be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (*e.g.*, a bovine, pig, sheep, rabbit, rat, mouse or other rodent) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at an embryonic (*e.g.*, one-cell) stage. Often all or substantially all of the somatic and germline cells of the transgenic animal have a copy of the transgene in their genome. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. Methods for generating transgenic animals, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Pat. Nos. 4,736,866 and 4,870,009. Typically, a GPRA and/or AAA1 nucleic acid is operably linked to a promoter and optionally an enhancer. The promoter and enhancer can be selected confer tissue specific expression. Transgenic animals that include a copy of a transgene introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of GPRA and/or AAA1 polypeptides. Such animals can be used as tester animals for reagents thought to confer protection from, for example, diseases related to GPRA and/or AAA1. Such a transgenic animal can be treated with agent and a reduced incidence of a characteristic of disease compared to untreated animals (*i.e.*, without the agent) bearing the transgene indicates a potential therapeutic intervention for the disease. Results from animal models can be confirmed in placebo-controlled clinical trials on patients with asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer.

Transgenic offspring are identified by demonstrating incorporation of the microinjected transgene into their genomes, preferably by preparing DNA from short sections of tail and analyzing by Southern blotting for presence of the transgene ("Tail Blots"). A preferred probe is a segment of a transgene fusion construct that is uniquely present in the transgene and not in the mouse genome. Alternatively, substitution of a natural sequence of codons in the transgene with a different sequence that still encodes the same peptide yields a unique region identifiable in DNA and RNA analysis. Transgenic "founder" mice identified in this fashion are bred with normal mice to yield heterozygotes, which are backcrossed to create a line of transgenic mice. Tail blots of each mouse from each generation are examined until the strain

is established and homozygous. Each successfully created founder mouse and its strain vary from other strains in the location and copy number of transgenes inserted into the mouse genome, and hence have widely varying levels of transgene expression or activity.

Knockout animals or cells can also be made by functionally disrupting an endogenous cognate form of one or more genes within the AST-1 locus (i.e., GPRA and/or AAA1). Functional disruption means that the transgenic animal is incapable of making a functional GPRA or AAA1 polypeptide and/or transcript encoding the same. Inactivation of a protein can be achieved by forming a transgene in which a cloned variant gene is inactivated by insertion of a positive selection marker within the coding region. See Capecchi, Science 244, 1288-1292 (1989). Inactivation of a transcript can be achieved by mutation within a promoter region. The transgene is then introduced into an embryonic stem cell, where it undergoes homologous recombination with an endogenous variant gene. Mice and other rodents are preferred animals. Such animals provide useful drug screening systems.

X. Methods of Screening for Drugs

The GPRA and AAA1 polypeptides and nucleic acids of the invention of the invention are useful for screening for agents that modulate (e.g., agonize or antagonize) their function (e.g., as a G-protein coupled receptor). Such agents can be useful as drugs in compensating for genetic variations affecting GPRA and/or AAA1 function. For example, an agonist of GPRA and/or AAA1 function is useful in a patient having a genetic variation that decreases endogenous GPRA function, and an antagonist is useful in a patient having a genetic variation that increases endogenous GPRA and/or AAA1 function. GPRA and AAA1 polypeptides can also be used to screen known drugs to determine whether the drug has an incidental effect on GPRA and/or AAA1 activity. A drug that has an incidental agonist activity should generally be avoided in patients having atypically high levels of IgE antibodies, and an antagonist should generally be avoided in immunosuppressed patients.

Agents can initially be screened in a binding assay with a GPRA or AAA1 polypeptides. A binding assay reduces the pool of initial agents to a subset that can be screened by a functional assay. Agents can be screened in cells transfected with nucleic acids encoding a GPRA and/or AAA1 polypeptide of the invention or in transgenic animals transfected with a GPRA and/or AAA1 nucleic acids and optionally and a reporter construct containing cAMP response element, a minimal promoter and a coding sequence for secreted alkaline

phosphatase. Methods for identifying agents that have a functional activity in transducing a signal through a cellular receptor are described by US 6,309,842. Agents for screening can be obtained by producing and screening large combinatorial libraries, as described above, or can be known drugs, or can be known ligands to related receptors to GPRA.

Agents can also be screened for capacity to inhibit or increase expression of GPRA and/or AAA1 nucleic acids. Suitable agents to screen include antisense polynucleotides, see *Antisense RNA and DNA*, (1988), D.A. Melton, Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY); Dagle et al., *Nucleic Acids Research*, 19:1805 (1991); Uhlmann et al., *Chem. Reviews*, 90:543-584 (1990), zinc finger proteins (see WO 00/00388 and EP.95908614.1), and siRNAs WO 99/32619, Elbashir, *EMBO J.* 20, 6877-6888 (2001) and Nykanen *et al.*, *Cell* 107, 309-321 (2001); WO 01/29058.

Agents having pharmacological activity evidenced by cellular or animal testing can be subject to clinical trials in patients suffering from asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer in comparison with a placebo.

XI. Methods of Treatment

The invention provides methods of treating asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer. These methods entail administering an effective amount of a GPRA and/or AAA1 polypeptide or an agent, e.g., an antibody, that modulates activity of a GPRA and/or AAA1 polypeptide. The agent can modulate that activity for example, by an effect on signal transduction through the GPRA polypeptide (usually a decrease) or by effecting expression of the GPRA and/or AAA1 polypeptide (also usually a decrease). An agent can be administered for prophylactic and/or therapeutic treatments. A therapeutic amount is an amount sufficient to remedy a disease state or symptoms, or otherwise prevent, hinder, retard, or reverse the progression of disease or any other undesirable symptoms in any way whatsoever. In prophylactic applications, an agent is administered to a patient susceptible to or otherwise at risk of a particular disease or infection. Hence, a "prophylactically effective" amount is an amount sufficient to prevent, hinder or retard a disease state or its symptoms. In either instance, the precise amount of compound contained in the composition depends on the patient's state of health and weight.

An appropriate dosage of the pharmaceutical composition is determined, for example, using animal studies (*e.g.*, mice, rats) are commonly used to determine the maximal tolerable dose of the bioactive agent per kilogram of weight. In general, at least one of the animal species tested is mammalian. The results from the animal studies can be extrapolated to determine doses for use in other species, such as humans for example.

The pharmaceutical compositions can be administered in a variety of different ways. Examples include administering a composition containing a pharmaceutically acceptable carrier via oral, intranasal, rectal, topical, intraperitoneal, intravenous, intramuscular, subcutaneous, subdermal, transdermal, intrathecal, and intracranial methods. The route of administration depends in part on the chemical composition of the active compound and any carriers.

The components of pharmaceutical compositions are preferably of high purity and are substantially free of potentially harmful contaminants (*e.g.*, at least National Food (NF) grade, generally at least analytical grade, and more typically at least pharmaceutical grade). To the extent that a given compound must be synthesized prior to use, the resulting product is typically substantially free of any potentially toxic agents, particularly any endotoxins, which may be present during the synthesis or purification process. Compositions for parental administration are also sterile, substantially isotonic and made under GMP conditions. Compositions for oral administration need not be sterile or substantially isotonic but are usually made under GMP conditions.

MATERIALS AND METHODS

Families were recruited in central eastern Finland in 1994 and 1996. The methods for recruitment, control for population stratification, and clinical evaluation have been described previously in detail (Kauppi *et al.* 1998; Laitinen *et al.* 1997). We used self-reported asthma as a sampling method. Altogether 253 families were recruited, two thirds of which were trios. Based on retrospective verification of the disease history and the results of diagnostic tests (spirometry, histamine or methacholine challenge test, expiratory peak flow measurements), 87% of the self-reported asthma patients were accepted as verified cases (Kauppi *et al.* 1998). Criteria for asthma were based on the recommendations of the American Thoracic Society (Dantzker, D.R. *et al.* 1987). In addition to the 86 multiplex pedigrees included already into

our previous genome scan, 103 nuclear families with full phase information (a total of 874 study individuals) were included to the association analysis without further selection.

Total serum IgE level was determined by Diagnostics CAP FEIA (Kabi Pharmacia, Sweden) in one batch. Allergy screening was done to all study individuals at the time of the study (Kauppi *et al.* 1998; Laitinen *et al.* 1997). Based on total serum IgE level the study individuals were divided into two groups: high IgE responders ($\text{IgE} > 100\text{kU/L}$) and low IgE responders ($\text{IgE} \leq 100\text{ kU/L}$).

EXAMPLES

EXAMPLE I

Construction Of The Physical Map

We have ordered the markers used in the analysis and estimated the physical distance between the markers using several publicly available sources as described previously by Polvi *et al.* (2002). Completed chromosome 7 sequence confirms the distance and order of our markers

(http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=Nucleotide&list_uids=22096428&dopt=GenBank).

EXAMPLE II

Microsatellite Discovery and Genotyping

To create a dense map of polymorphic markers spread evenly across the linkage region, we screened the publicly available genomic sequences for potentially polymorphic tandem repeats (Polvi *et al.* 2002). Eighty microsatellites were genotyped in the whole data set (874 samples) giving the average marker density of 220kb (range from 904bp to 790kb) across the linkage region. All microsatellite markers and small insertions and deletions (Table 1C) found in the critical region were genotyped using fluorescently labeled primers in gel electrophoresis on the ABI377 sequencer as described previously (Polvi *et al.* 2002 and <http://www.genome.helsinki.fi/eng/research/asthma/detail.htm>).

EXAMPLE III

Discovery of Genetic Variation in the Critical Region by Direct Sequencing

To identify all sequence polymorphisms in the critical region, we generated a 129 kb reference sequence by assembling the sequence (genomic clones AC005680, AC005492, AC005174, AC005862.1, and AC005853.1) from the Human Genome Project (in the contig NT_000380 from the position 509,783 to the position 638,799). SNP discovery by re-sequencing was done in four selected patients and in three control individuals who were homozygous across the region and for the flanking markers. Repeat regions such as SINE, LINE, LTR, MER1 and MER2 elements covered 60% of the sequence (a total of 129,017 bp). PCR assays were designed to be ~650 bp in length with 100 bp overlap with adjacent assays. Repeat regions were re-sequenced when it was possible to design primers in unique sequence within the range mentioned above.

PCR assays were carried out in 20µl volumes containing 20ng of genomic DNA, 1x PCR Buffer II, 0.1mM dNTPs, 2.5 mM MgCl₂, 0.1µM primer mix, and 0.5 U of DNA polymerase (AmpliTaqGold, Applied Biosystems). The samples were denatured for 10 min at 94°C, followed by 35 cycles each of 30s at 94 C, 30 s at 58 C, and 30 s at 72 °C. Elongation was performed for 10 min at 72°C. Purified PCR fragments (Quickstep 2 PCR purification Kit, Edge BioSystems, MD) were sequenced in both directions using ABI Prism3100 sequencer and dye-terminator chemistry. We assembled the sequence reads using the Gap4 program.

EXAMPLE IV

SNP Genotyping

SNP genotyping was done using two different methods: single base pair extension (SBE) or using altered restriction sites. SBE was done using the chemistry of Molecular Dynamics according to the suggestion made by the manufacturer on a Megabase 1000 sequencer (Molecular Dynamics, CA) using the primers listed in the Table 1A. Allele calling was performed by using the MegaBACE SNP Profiler software (Molecular Dynamics).

Twenty-nine of the SNPs were genotyped using different restriction enzyme digestions. All primers, restriction enzymes, and lengths of the digestion fragments of corresponding allele used in the genotyping are given in Table 1B. If the SNP did not produce a natural site for

altered restriction, mutations were induced in PCR-primers (shown in capital letters in primer sequence). To improve allele calling by growing size difference between alleles, in some primers plasmid sequence tail of 20-30bp was added (shown in bold in the primer sequence). Altered restriction sites of the PCR products stained with ethidium bromide and visualized on agarose gels in UV light were called manually by two independent observers. All the markers were in Hardy-Weinberg equilibrium and observed Mendel errors were less than 0.1%.

EXAMPLE V

Haplotype Association Analysis

The haplotype analysis was done using Haplotype Pattern Mining (HPM) (Toivonen *et al.* 2000) program. For haplotyping, large pedigrees were divided into trios using an in-house computer program. The program identifies the maximum number of trios that are not overlapping and in which one or two members were affected (not both parents). Trios that included members who had not been genotyped or members with unknown phenotype were excluded. In phase three of hierarchical LD mapping, high density genotyping of the critical region was done in the trios informative for the association analysis (a total of 132 trios, 396 study individuals, and revealing 304 unrelated affected and 220 control chromosomes). Haplotyping was done within each trio and four independent chromosomes were obtained from each trio. In case of ambiguities (missing genotype data, identical heterozygotic genotypes in all of the family members, or Mendel errors), the alleles were discarded. If the child was affected, the transmitted chromosomes were considered disease associated and the non-transmitted chromosomes as controls. If one of the parents was affected, his/her chromosomes were considered disease associated and the spouse's chromosomes as controls. If both the parent and the child were affected, only the non-transmitted chromosome of an unaffected parent was considered as the control and the other three as disease associated. These haplotypes were used as input for HPM.

To estimate the overall significance of detected association, accounting for simultaneous testing of multiple markers, we used a second level permutation test by performing nested permutation tests described in detail elsewhere (Sevon *et al.* 2001a and b).

RESULTS

1. Haplotype Association

Our genotyping effort searching for a shared haplotype among individuals who have high total IgE serum level was done in three phases using a hierarchical approach. Based on the initial observation on potential association, the marker map was made denser in the regions of primary interest. In the haplotype analysis we used the HPM algorithm that is known to be powerful in locating a disease-causing gene when one is known to exist (Toivonen *et al.* 2000). By allowing gaps in the haplotype patterns, the algorithm is robust for genotyping errors, marker mutations, unrecognized recombinations and missing data. The data set consisted of 86 multiplex pedigrees that were included also in the previous linkage study and additional 103 nuclear families recruited solely for the association study. These families revealed 304 unrelated high IgE associated and 220 control chromosomes which were then used as the input data for the HPM analysis.

First we built a marker map across the whole linkage region with 80 microsatellite markers (Fig. 1). In the first phase of haplotype association analysis the marker density was approximately 220kb across the linkage region. The best associations (chi square ≥ 7.4) formed only one cluster of haplotypes located between the markers D7S690–NM13 (~ 3.21 cM). In the second phase we added 6 new microsatellites in between D7S690 and NM13. The associated haplotypes (chi square ≥ 8.9) clustered now at the centromeric end of the region, between markers G42099 and NM13 (301kb) (Fig. 2). HPM algorithm was used to evaluate the overall haplotype distribution among the high IgE associated compared to that among control chromosomes. A permutation test incorporated to the HPM computer package was used to evaluate the statistical significance of our finding by comparing the overall haplotype distribution among the high IgE associated and control chromosomes. Using chi square 7.0 as the threshold for association, 7 markers as the maximum length of the haplotype, and allowing one gap in the haplotype patterns, the best associations were computed for three adjacent markers G42102, D7S497, and G4296 (marker-wise P values 0.05-0.06 based on 10 000 simulations).

Fine mapping was then further continued by adding 49 new markers between G42099 and NM13 giving an average marker density of 6kb. By using the HPM algorithm, all associated haplotype (chi square > 13) patterns spanned between SNP509783 and SNP638799 (129,017 bp).

To study if the overall haplotype distribution in the locus differs in high IgE associated compared to that in control chromosomes, we used again a permutation test. HPM analysis was done using the following parameters: maximum pattern length 40 markers, one gap allowed for missing data and possible errors, and chi-square threshold for the association 13.0. The observed scores (=number of qualified haplotype patterns spanning across the marker) for associated haplotypes varied from 0 to 40. Permutation test showed statistically biased haplotype distribution for high IgE. The marker-wise P values ≤ 0.005 were observed for all eleven of the markers between SNP531632 and SNP563930 and the marker wise P values ≤ 0.01 for all twenty-eight markers between NM51 and SNP617392 based on 10 000 simulations. To estimate the effect of simultaneous testing of multiple markers, we used a second level permutation test, yielding a corrected P value of 0.03 (Sevon 2001a and b).

2. Genetic Variation in the Critical Region

Based on haplotype analysis we had determined the critical region in between the markers SNP509783 and SNP638799 (Fig. 2). SNP genotyping resulted in the identification of seven haplotypes H1-H7 that together explain 94.5% of the genetic variation detected at the AST-1 locus. To identify all potentially causative genetic variations on the risk haplotype, we sequenced four patients who carried haplotypes H2/H2, H4/H4, H5/H5, and H7/H7, respectively, and were homozygous across the AST1-locus region. The patients' sequences polymorphisms were compared with the reference sequence (SEQ ID NO:1)

(http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=nucleotide&list_uids=7382309&dopt=GenBank, human genomic contig NT_000380) and to three population based controls homozygous for the same markers, but with different haplotypes H1/H1, H3/H3, and H6/H6, respectively. The differences are highlighted via SEQ ID NO:1. Genetic variation including small deletions and insertions (≤ 6 base pairs), variation in the length of microsatellite repeats, and multiple SNPs are presented in Table 3 that can be used as genetic markers of the risk haplotypes. Tables 13 and 14 show the SNPs unique for each haplotype and unique for certain haplotype combinations, respectively. In diagnostic testing these markers identify the haplotypes and the haplotype combinations of an individual without phase information.

Example VI: Identification of the GPRA Gene

A. Materials and Methods

1. Exon Prediction

Exon predictions were performed by GENSCAN (<http://genes.mit.edu/GENSCAN.html>) and software using genomic clones from AC005493.1 to AC005826.1 (690 kb) (Polvi *et al.* 2002) as one block. Predictions were performed for both unmasked and masked sequence. In the latter, the repetitive sequences were masked using the Repeat Masker Service (<http://repeatmasker.genome.washington.edu/>).

2. Reverse Transcriptase PCR

The exon-specific primers were designed based on exon predictions (Table 4, Fig. 3). First strand cDNAs were synthesized from 1 µg of commercially available poly A⁺ RNAs from human brain (Invitrogen), lung, testis, placenta, and thymus (BD Clontech) by the SMART RACE cDNA Amplification Kit (BD Clontech, Palo Alto, CA) according to manufacturer's instructions. Alternatively, Marathon-Ready cDNA (Clontech) or cDNA from Multiple Tissue cDNA Panels I and II (Clontech) were used as templates in some PCR amplifications. All the PCR amplifications were performed in 20-50 µl volumes using 2.0-5.0 µl cDNA as template, 2 mM MgCl₂, 0.2 mM dNTPs (Finnzymes), 0.2 mM of each primer, 0.15-0.4 U AmpliTaqGold, and 1x PCR Gold buffer (Applied Biosciences) under the following conditions: 94°C for 10 min, 35-40 cycles of 94 °C for 30 s, 58 °C –68 °C for 30 s, 72 °C for 1 min 40 s to 2 min followed by 72 °C for 10 min. PCR products were analyzed on 1 % agarose gel and extracted from the gel using QIAquick Gel Extraction Kit (Qiagen).

3. Rapid Amplification of cDNA Ends (RACE-PCR)

To generate 3' and 5' cDNA ends, rapid amplification of cDNA ends (SMART RACE) was performed using human testis cDNA and Human Marathon-Ready Fetal Thymus cDNA (Clontech) according to the manufacturer's protocol for the SMART RACE cDNA Amplification Kit (BD Clontech). RACE-PCR products were cloned using pGEM-T Easy Vector system (Promega) or TOPO TA Cloning kit (Invitrogen) and plasmid DNA was purified using QIAprep Spin Miniprep Kit (Qiagen).

The purified RT-PCR products and the cloned RACE-PCR products were verified by automated sequencing with dye-terminator chemistry (ABI Prism3100, Applied Biosystems, Inc).

4. Cloning of the Full-Length cDNAs of GPRA

Nested PCR amplification was used in cloning of the full length cDNAs for *GPRA* splice variants A and B. The first round of PCR for variant A was done with the primer pair JEGE1F1 and JEGE9aR1, and for variant B with JEGE1F1 and JEGE9bR1 (Table 4, Fig. 3). The following inner primers were used in the re-amplification: JEGE1F2 and JEGE9aR2 for variant A; JEGE1F2 and JEGE9bR2 for variant B, respectively. Full length C variant was amplified with the primer pair JEGE1F2 and JEGExR1.

Primary PCR amplifications of variants A and B were performed in 25 μ l volumes using 2.5 μ l Human Brain Marathon-Ready cDNA (Clontech) as template, 1x DyNAzyme EXT buffer (containing 1.5 mM $MgCl_2$), 0.2 mM dNTPs (Finnzymes), 0.52 μ M of each primer, 5% DMSO and 0.5 U DyNAzyme EXT polymerase (Finnzymes) under the following conditions: 94°C for 4 min, 38 cycles of 94°C for 30 s, 65°C for 30 s, 72°C for 1 min followed by final extension of 72°C for 10 min. The aliquot of primary PCR product was re-amplified by 30 PCR cycles under the same conditions as above. PCR amplification of variant C was performed in the above conditions. NCI-H358 cDNA was used as template.

PCR products were cloned into pCR 2.1 TOPO -vector using TOPO TA cloning kit (Invitrogen) according to manufacturer's instructions and plasmid DNA were purified using QIAprep Spin Miniprep Kit (Qiagen). The cloned RT-PCR products were verified by automated sequencing with dye-terminator chemistry (MegaBACE 1000, Molecular Dynamics).

5. Cell Culture and Isolation of Poly A⁺ RNA

In addition to commercially available poly A⁺ RNAs and Marathon-Ready cDNAs, human lung epithelial carcinoma cell line NCI-H358 (ATCC) was cultured as mRNA source for expression studies. Cells were cultured in RPMI 1640 medium (Gibco BRL) supplemented with 1 mM sodium pyruvate (Gibco BRL), 10% FCS (Biological Industries), and 1% Penicillin/Streptomycin (GibcoBRL). Poly A⁺ RNA was isolated by Dynabeads mRNA DIRECT Kit (Dynal) according to the manufacturer's instructions.

6. Northern Blot Hybridization

The GPRA C specific (comprising the nucleotides from 58 to 527 of the *GPRA* C variant) 470 bp probe was generated by RT-PCR using human testis cDNA as template. The probe was radiolabelled with α [^{32}P]-dCTP using RediPrime Kit (Amersham Biosciences) according to the manufacturer's instructions. A Human Multiple Tissue 8-lane Northern blot (BD Clontech) was prehybridized in ExpressHyb solution (BD Clontech) for 1 h at 68 °C followed by hybridization with the specific probe for 1 h 15 min at 68 °C. Herring sperm DNA (100 $\mu\text{g/ml}$) was used as the blocking reagent. The filter was washed with 2 x SSC and 0.05 % SDS at room temperature and then exposed to X-ray film at -20 °C for 6 days.

7. Western Blot and Immunohistochemistry

Two specific antibodies against the alternative carboxy terminals of *GPRA* were raised by immunizing rabbits with the following peptides: CREQRSQDSRMTFRERTER (corresponding to the nucleotides 1148-1205 of the variant A) and CPQRENWKGTWPGVPSWALPR (corresponding to the nucleotides 1196-1259 of the variant B of *GPRA*). GPRA A peptide synthesis and antibody production were purchased from Sigma-Genosys Ltd (London Road, Pampisford, Cambridge). GPRA B peptide synthesis was purchased from the University of Helsinki and antibody production from the University of Oulu. A total of 6 immunizations were performed at 2 weeks intervals using the total of 2 mg of KLH-conjugated peptide purified by gel filtration. GPRA antibodies were purified from whole serum by affinity chromatography with the peptide coupled to iodoacetyl on a crosslinked agarose support according to the manufacturer's (Pierce, Meridian Road, Rockford IL, USA) instructions.

For western blot analysis, human tissue lysates from spleen, skeletal muscle, uterine muscle, colon muscle, kidney, colon epithelium, testes, and prostate were obtained by mechanically homogenizing the frozen tissue samples in 10 mM Tris HCl 100mM NaCl 2 % Triton X-100 buffer with proteinase inhibitors. 50 μg of the protein lysates were run on reducing 12.5 % SDS-PAGE gels and electroblotted to the PVDF membrane according to standard procedures. Nonspecific protein binding was prevented by incubating the membrane with 5 % milk in 0.1 % Tween 20 in TBS (TBST). Antigenic sites were revealed by incubating the membrane with the anti-GPRA C-terminal antibodies or pre-immune serum followed by the alkaline phosphatase-conjugated goat anti-rabbit secondary antibody (Jackson ImmunoResearch Laboratories Inc., West Baltimore Pike West Grove, PA, USA) in 5 % milk in 0.1 % TBST.

The color reaction was revealed by the NBT/BCIP method (Pierce). Negative controls did not show specific reactivity.

Formalin fixed, paraffin-embedded specimens of normal adult human bronchus, skin and colon, and human normal tissue array slides (MaxArray, Zymed Laboratories Inc., CA) containing 30 different tissues were used for immunohistochemistry. For pre-treatment, slides were deparaffinized by xylene-treatment followed by decreasing alcohol series. The slides were heated in microwave oven in 10 mM citrate buffer, pH 6.0 for 5 minutes.

Immunohistochemical analyses were performed using the ABC method (Vectastain Elite ABC kit, Vector Laboratories, Burlingame, California). Omission of primary antibody and staining with the nonimmunized sera was used as negative control for parallel sections. Neither of these controls showed any immunoreactivity.

8. Transient transfections and Elisa

Cos-1 cells were cultured in Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum, 1% PS and 5% non-essential amino acids. Cos-1 cells were transiently transfected with pCMV-myc- GPRA A, B, C, D, E, F or Bshort separately using Fugene6 transfection reagent (Roche) following the manufacturer's protocol. After 24 hours, the transfected cells from one well of 6 well plate were divided into 16 wells of 96 well plate. After 48 hours of transfection, the cells were fixed with 3.5 % PFA. Half of the cells were permeabilized with 0.5% TX-100 for 10-15 min, blocked with TBS containing 2% milk and 1% goat normal serum at 37 °C for 30 min, incubated with 1:1000 dilution of myc- specific primary antibody (Babco) for 1 h at 37 °C, washed three times with TBS and thereafter incubated with a dilution of 1:2000 of HRP-conjugated anti-mouse IgG antibody for 30 min at room temperature and washed three times with TBS. TMB-substrate (Sigma) was added to cells and the reaction was let to proceed for 3 to 6 minutes after which the reaction was stopped by adding 1.5 M HCL. Absorbance was measured at 450 nm.

9. Computational Protein Sequence Analysis

The transmembrane topology and putative N-glycosylation sites of *GPRA* protein were predicted using PredictProtein (http://cubic.bioc.columbia.edu/predictprotein/submit_def.html#top) and TMPred (http://www.ch.embnet.org/software/TMPRED_form.html) softwares. In protein homology comparisons the BlastP

(<http://www.ncbi.nlm.nih.gov/blast/>) and T-Coffee (<http://www.ch.embnet.org/software/TCoffee.html>) softwares were used.

10. Family Collection

Genotyping of *GPRA* was done in a family collection recruited in central eastern Finland. The methods for recruitment, control for population stratification, and clinical evaluation have been described previously in detail (Kauppi *et al.* 1998; Laitinen *et al.* 1997). We used self-reported asthma as a sampling method. Total serum IgE level was determined by Diagnostics CAP FEIA (Kabi Pharmacia, Sweden) in one batch for all the participants. Based on total serum IgE level the study individuals were divided into two groups: high IgE responders (IgE > 100kU/L) and low IgE responders (IgE ≤ 100 kU/L).

Altogether 253 families were recruited, two thirds of which were trios. Based on retrospective verification of the disease history and the results of diagnostic tests (spirometry, histamine or methacholine challenge test, expiratory peak flow measurements), 87% of the self-reported asthma patients were accepted as verified cases (Kauppi *et al.* 1998). Criteria for asthma were based on the recommendations of the American Thoracic Society (Dantzker, D.R. *et al.* 1987). 86 large pedigrees were included into our previous genome scan (Laitinen *et al.* 2001). For the association study, those pedigrees were divided into trios using a in-house computer program. Non-overlapping trios with full phase and phenotype information and additional 103 trios were included to the association analysis without further selection.

11. Screening for Exon Polymorphisms

Genomic DNA of four asthma patients and one healthy control from above mentioned study group were screened for sequence variation with 12 primer pairs covering all the verified exons of *GPRA* (Table 5). All the patients were homozygous for microsatellites G42099, G42100, D7S497, and G42097 (from the NT_000380 position 476,213 to 644,425, a total of 168 kb) and expressed the susceptibility haplotype (Patent 1). The control individual showed homozygosity for the same markers, but to a different, non-associating haplotype.

PCR assays were carried out in 20 µl volumes containing 20 ng of genomic DNA, 0.1mM dNTPs (Finnzymes), 2.5 mM MgCl₂ (Applied Biosystems), 0.1µM primer mix, and 0.5 U of DNA polymerase, and 1x PCR Buffer II (AmpliTaQGold, Applied Biosystems). The samples

were denatured for 10 min at 94 °C, followed by 35 cycles each of 30s at 94 °C, 30 s at 58 °C, and 30 s at 72 °C. Elongation was performed for 10 min at 72 °C. Purified PCR fragments (Quickstep 2 PCR purification Kit, Edge BioSystems) were sequenced from both directions using ABI Prim3100 (Applied Biosystems) sequencer and dye-terminator chemistry. We assembled forward and reverse sequence reads using the Gap4 program (Staden Package software).

12. Genotyping

Genotyping of five exonic SNPs in *GPR4* was done using two different methods. One marker was genotyped using single base pair extension (SBE) with the chemistry of Molecular Dynamics on a Megabase 1000 sequencer (Molecular Dynamics) according to the suggestion made by the manufacturer (Table 6A). Allele calling was performed by using the MegaBACE SNP Profiler software (Molecular Dynamics).

Four of the SNPs were genotyped using different restriction enzyme digestions. All primers, restriction enzymes, and lengths of the digestion fragments of corresponding allele used in the genotyping are given in Table 6B. If the SNP did not produce a natural site for altered restriction, mutations were induced in PCR-primers. To improve allele calling by growing size difference between alleles, in one primer plasmid sequence tail was added. Altered restriction sites of the PCR products were visualized on ethidium bromide stained agarose gels in UV light and called manually by two independent observers. All the markers were in Hardy-Weinberg equilibrium and observed Mendel errors were less than 0.1%.

13. Haplotype Association Analysis

The haplotype analysis was done using Haplotype Pattern Mining (HPM) (Toivonen *et al.* 2000) program. The data set provided a total of 132 informative trios, 396 study individuals, and revealing 304 unrelated affected and 220 control chromosomes. Haplotyping was done within each trio and four independent chromosomes were obtained from each trio. In case of ambiguities (missing genotype data, identical heterozygotic genotypes in all of the family members, or Mendel errors), the alleles were discarded. If the child was affected, the transmitted chromosomes were considered disease associated and the non-transmitted chromosomes as controls. If one of the parents was affected, his/her chromosomes were considered disease associated and the spouse's chromosomes as controls. If both the parent

and the child were affected, only the non-transmitted chromosome of an unaffected parent was considered as the control and the other three as disease associated. These haplotypes were used as input for HPM.

RESULTS

1. Characterization GPRA at RNA And Protein Level

A. Genomic Structure of Different Splice Variants of GPRA

The full-length cDNA sequence of *GPRA* was assembled by using RT-PCR with primers designed for predicted exons and by using RACE-PCR to generate the 5' and 3' ends of the gene. The genomic location of the primers is shown in the Fig. 3 and sequence in the Table 4. Nested PCR amplification was used to produce full length cDNAs of different splice variants in one fragment. Using different (Marathon-Ready Brain cDNAs, RNA extracted from NCI-H358 cell line) template three splice variants were identified repeatedly (A, B_{long}, and C variants in the Fig. 3 and 4). All variants possess the same initiation site, but used alternative exons to encode the 3' end of the gene (exons 9A, 9B and 2B, respectively). The sequence flanking the putative ATG translation initiation site (GCCATGC) contains the -3 purine, but not the +4 guanine residue, of the Kozak consensus sequence. In two of the variants, the open reading frame was distributed across 9 exons producing cDNAs of 1116 bp and 1134 bp in size (variants A and B_{long}, Fig. 5A and 5B) and encoding 371 and 377 amino-acid proteins, respectively. The C variant is a shorter transcript of *GPRA* including only the exons E1, E2a and E2b (Fig. 5C). *GPRA* spanned 0.2 Mb of the genomic contig NT_000380. The second intron is the largest, comprising of 93.8 kb.

When nested PCR specific for the B variant was applied in the cloning process a new splice variant was observed (B_{short} in Figs. 4B and 5B). A 33 bp deletion was observed in the 5' prime end of the exon 3.

B. Expression Profiling of GPRA Using RT-PCR and Northern Blot Hybridization

RT-PCR analysis showed that different variants of *GPRA* were expressed in several human tissues. These tissues include testis, brain, pituitary gland, placenta, lung, heart, fetal thymus, and fetal heart. In addition to commercially available cDNAs, we analyzed the *GPRA* expression profile in NCI-H358 cell line that represents broncho-epithelial origin. Variants A (JEG5F1 and Vau8R2), B (JEG5F1 and Vau1000R1) and C (JEG5F1 and JEGEXR1)

specific primer pairs were used in amplifications (Fig. 8, Table 4). All forms were expressed and using A variant specific primers, we found several transcripts. PCR fragments were cloned and sequenced. Sequence verification revealed variants A, D, E, and F (Figs. 4 and 5). Northern blot hybridization was done using a 470 bp cDNA probe comprising exons E1, E2A, and E2B (Fig. 3). On Human 8-Lane MTN blot, the probe detected multiple transcripts sized approximately 6.5 kb, 6.0 kb, 1.8 kb, and 1.0 kb as shown for placental tissue in Fig. 9.

C. The Predicted Structure of the GPRA Protein

Both TMpred and PredictProtein softwares predicted that the variants A and B_{long} encode a 7TM protein with an extracellular N-terminus (approximately 50 amino acids) and intracellular C-terminus (Figs. 6A-6F). Their structure shared the common features of GPR family A with 16 conserved amino acids in the TM, loop, and C-terminal regions. These include among others two conserved Cys residues in exoloops 1 and 2 that potentially form a disulfide bridge; the Asp-Arg-Tyr sequence (DRY motif) in the proximity of TM3; Asn-Pro-X-X-Tyr motif in TM7; and a Cys residue in the C-terminal region (Figs. 6A-6F). The putative N-glycosylation site Asn⁴ was predicted by PredictProtein. All the other variants lacked the 7TM structure. For variant B_{short} a 6TM structure was predicted (Fig. 6B). Variant C encodes 94 amino acid protein that has potentially only one TM region (Fig. 6C). For the variants D and E, deletions of the exons 3 or 4 caused a shift in the ORF, presumably producing truncated forms of the protein. When both exons were deleted (variant F), the ORF remained intact, but only five-transmembrane regions were predicted (Fig. 6F).

D. Sequence Searches

Determined by BlastP comparisons, the GPRA protein shows 31 % amino acid identity to human vasopressin receptor 1B, 28 % identity to human vasopressin receptor 2, 32 % identity to human oxytocin receptor, and 43 % amino acid sequence identity to gene product CG6111 of *Drosophila melanogaster* which is considered as an orthology to human vasopressin/oxytocin receptor family (Park *et al.* 2002). The best sequence identity between proteins was found in TM regions and in the first exoloop. Based on T Coffee homology comparisons, GPRA showed 20% amino acid identity to the bovine rhodopsin receptor. Sequence comparisons between GPRA and mouse ESTs BB638128, BB632343, BB625809, BB228269, and publicly available mouse genomic sequence suggest the existence of a mouse orthology.

2. Expression Profiling of GPRA Variants A and B Using Western Blot Analysis and Immunohistochemistry

Immunostaining of normal adult human tissue samples with GPRA A variant specific antibody showed strongest expression in smooth muscle cells (SMC), such as SMC layer in bronchial and arterial walls in human lung and colon shown in Fig. 10A and E. In alveolar wall and alveolar macrophages of lung (Fig. 10C) intense staining of *GPRA A* was detected. In colon epithelium (Fig. 10D), strong basal staining was observed, whereas in the bronchial epithelium (Fig. 10A) and keratinocytes (Fig. 10F), only mild/weak immunostaining was detected.

Western blot analysis with GPRA A variant specific antibody revealed four intensive bands corresponding to molecular weights of approximately 50, 44, 42 and 40 kDa (Fig. 12A). In skeletal muscle only 50 kDa band was detectable whereas uterine muscle, colon epithelium and prostate showed similar expression patterns at 50, 44, and 40 kDa. The most intensive GPRA A expression was recorded in colon muscle with the major band at 42 kDa. Weak or nonexistent bands were found in spleen, kidney, and testis.

Immunostaining with GPRA B variant specific antibody revealed most intense expression of the protein in the epithelium of several mucosal tissues including bronchus, small intestine, and colon as shown in Figs. 11A, D, E. Contrary to the A variant, the expression of the B variant was strongest in the apical surface of the epithelium (Figs. 11A, D, E, and F). Alveolar walls and alveolar macrophages (Fig. 11C) showed strong and SMCs in several tissues mild immunostaining of the variant B.

With GPRA B variant specific antibody, two sharp polypeptide bands corresponding to molecular weights of approximately 39 and 25 kDa were revealed by Western blot analysis with the strongest expression in kidney (Fig. 12B). The specificity of the antibody was confirmed by the blocking experiment with 10x molar excess of the peptide used in immunization.

Cos-1 cells were transiently transfected with pCMV-myc GPRA A, B, Bshort, C, D, E, or F in order to study translocation of different GPRA variants *in vitro*. After 48 hours, cells were fixed and studied by cell based ELISA assay using myc antibody. 71% of recombinant GPRA

A and 52% of recombinant GPRA B were translocated to the plasma membrane while all the other variants were located in cytoplasm (Table 15).

3. Characterization Of GPRA as a Genetic Regulator of Asthma Related Traits

A. Detection of GPRA Polymorphisms Associated to Asthma Related Traits

To detect sequence variations in *GPRA* associated to asthma related traits, we sequenced its verified exons and exon-intron boundaries in four patients homozygous for the haplotype significantly associated with high serum total IgE level among the Finnish asthma families. Comparison between patients and one control individual from the same data set were made to the reference sequence (genomic contig NT_000380) in the public data base.

A total of 14 SNPs were found distributed in seven exons of *GPRA* (Table 7). Four of the SNPs were predicted to cause a non-conservative amino acid change: Asn>Ile located in the first extracellular loop, Ser>Arg in the third cytoloop, Gln>Arg and Thr>Ile in the C termini, one in each alternative splice variant. Six of the SNPs were located in the 3' UTRs of the A, B, and C variants of the gene.

B. Association Analysis

We have previously defined a susceptibility haplotype that significantly associates with asthma related phenotypes (best marker-wise P value based on 10,000 permutations 0.001) (Fig. 3, gray area). The region covers the genomic region between *GPRA* exons 2b and 5. We replicated the association analysis with five SNPs genotyped in the exons of *GPRA* in the same data set (304 disease associated and 220 control chromosomes). Best haplotype associations of *GPRA* for high serum IgE level when one gap was allowed in the haplotype patterns are shown in Table 8. The best observed associations reached χ^2 values of 5.7-10.3. Only one of markers (SNP591694) was part of the previously determined susceptibility haplotype and the same amino acid change, 107Ile, showed also significant association to high IgE level together with other *GPRA* SNPs.

To study whether the overall haplotype distribution in *GPRA* differs in high IgE associated compared to that in control chromosomes, we used a permutation test. HPM analysis was done using the following parameters: maximum pattern length 5 markers, one gap allowed for missing data and possible errors, and chi-square threshold for the association >5.0. The

observed scores (=number of qualified haplotype patterns spanning across the marker) for associated haplotypes varied from 6 to 14. Permutation test showed statistically biased haplotype distribution for high IgE. Based on 10,000 simulations, the best marker-wise P value ≤ 0.01 was observed for a silent SNP (in position 640,764) in exon 5. P ≤ 0.02 was observed for three markers in the middle of the haplotype and P < 0.04 for all the markers in the haplotype.

Example VII

This example describes a second gene occurring partially within the AST-1 locus and overlapping the GPRA gene.

A. Materials and Methods

1. Exon Prediction

Exon predictions were performed by GENSCAN (<http://genes.mit.edu/GENSCAN.html>) software using genomic clone NT_000380 (Polvi *et al.* 2002). Predictions were performed for both unmasked and masked sequence. In the latter, the repetitive sequences were masked using the Repeat Masker Service (<http://repeatmasker.genome.washington.edu/>).

2. Reverse Transcriptase PCR

The exon-specific primers were designed based on exon predictions (Table9, Figure13). First strand cDNAs were synthesized from 1 μ g of commercially available poly A⁺ RNAs from human lung, testis, kidney and fetal liver (BD Clontech) by the SMART RACE cDNA Amplification Kit (BD Clontech, Palo Alto, CA) according to manufacturer's instructions. Alternatively, Marathon-Ready cDNA (Clontech) or cDNA from Multiple Tissue cDNA Panels I and II (Clontech) were used as templates in some PCR amplifications.

All the PCR amplifications were performed in 25-50 μ l volumes using 2.0-5.0 μ l cDNA as template, 2 mM MgCl₂, 0.2 mM dNTPs (Finnzymes), 0.2 mM of each primer, 0.15-0.4 U AmpliTaqGold, and 1x PCR Gold buffer (Applied Biosciences) under the following conditions: 94°C for 5 min, 35-40 cycles of 94 °C for 30 s, 58 °C – 62 °C for 40 s, 72 °C for 1 min 30 s followed by 72 °C for 10 min. PCR products were analyzed on 1 % agarose gel and extracted from the gel using QIAquick Gel Extraction Kit (Qiagen). Purified PCR

products were analyzed by automated sequencing with dye-terminator chemistry (Megabase 1000 sequencer, Molecular Dynamics).

3. Cell Culture and Isolation of Poly A⁺ RNA

In addition to commercially available poly A⁺ RNAs, RNA from Epstein Barr virus infected lymphoblast cell lines of the patients who were homozygous, heterozygous, or non-carriers of AST1 were cultured for expression studies. Lymphoblasts were cultured in RPMI 1640 medium (Gibco BRL) supplemented with 1 mM sodium pyruvate (Gibco BRL), 10% FCS (Biological Industries), and 1% Penicillin/Streptomycin (GibcoBRL). Poly A⁺ RNA was isolated by Dynabeads mRNA DIRECT Kit (Dyna) according to the manufacturer's instructions.

4. Northern Blot Hybridization

The AAA1 specific probe (mixture of the variants I, III, IV, VII, and X) was generated by RT-PCR using human lung, testis, kidney and fetal liver poly A⁺ RNAs as templates. The probe was radiolabelled with α [³²P]-dCTP using RediPrime Kit (Amersham Biosciences) according to the manufacturer's instructions. A Human Multiple Tissue 12-lane Northern blot, Fetal Multiple Tissue Northern blot, and Human Multiple Tissue Expression Array 2 (BD Clontech) were prehybridized in ExpressHyb solution (BD Clontech) for 1 h at 68 °C followed by hybridization with the specific probe for 2-5 h at 68 °C. Herring sperm DNA (100 μ g/ml) was used as the blocking reagent. Filters were washed with 2 x SSC and 0.05 % SDS at room temperature and exposed to X-ray film at -20 °C for 1 - 5 days.

5. Genotyping

Genotyping of four SNPs that are located either in the coding region or near the exon-intron boundaries of AAA1 was done 1) using altered restriction sites (SNP_538567 or SNP_574953, Table 1B) or 2) single base pair extension (SBE) with the chemistry of Molecular Dynamics on a Megabase 1000 sequencer (Molecular Dynamics) according to the suggestion made by the manufacturer (Table 10). Allele calling was performed by using the MegaBACE SNP Profiler software (Molecular Dynamics).

All the markers were in Hardy-Weinberg equilibrium and observed Mendel errors were less than 0.1%.

6. In vitro translation of the AAA1 and characterization of AAA1 antibody

In order to investigate whether AAA1 is translated to any polypeptide, *in vitro* translation experiments were performed (Figure 21). Capped RNAs of AAA1 gene variants defined by SEQ ID NOS: 16 and 22 were transcribed from DNA constructs with the aid of T7 RNA polymerases (mMESSAGE mMACHINE system, Ambion, USA). Translation was performed with rabbit reticulocyte lysate translation machinery (Riboprobe *in vitro* translation system, Promega) in the presence of S³⁵-labelled methionine in the reaction mixture. The *Xenopus* elongation factor α (pTRI-Xef) DNA template was used as a positive control for transcription and translation. In negative control, water was used instead of DNA. The translated polypeptides were detected by autoradiography after Tris-Tricine SDS-PAGE.

B. Results

1. Characterization of AAA1 at RNA Level

A. Genomic Structure of Different Splice Variants of AAA1

The human AAA1 gene spans 520 kb of the genomic contig NT_000380 (nucleotides 163615–684776) and it is divided into 18 exons (Figure 13 and Table 11). The cDNA sequences of AAA1 splice variants were assembled by using RT-PCR with primers designed for predicted exons. The genomic location of the primers is shown in the Figure 13 and the sequence in the Table 9. Using cDNAs from human lung, kidney, testis and fetal liver as templates, twelve splice variants were identified repeatedly (variants I–XII in the Figure 14). All variants share exon 6, but use alternative exons to encode the 5' and 3' ends of the transcript. In six variants out of ten, the sequence flanking the putative ATG translation initiation site (GCCATGC) contains the –3 purine, but not the +4 guanine residue, of the Kozak consensus sequence.

B. Expression Profiling of AAA1 Using RT-PCR and Northern Blot Hybridization

Northern blot analysis of a multiple tissue expression array showed that AAA1 is expressed in several human tissues (Figure 16). Strong expression was seen in testis, brain, placenta, lung, heart, skeletal muscle, kidney, liver, fetal liver, and fetal lung. In multiple tissue northern blots, two main transcripts (2.4 and 7.5 kb in size) were detected (Figure 17). Additionally, all sample lanes showed extensive label streaming which can be an indication of several low-abundant transcripts with different sizes.

By RT-PCR exceptionally strong splicing and tissue specific differences in AAA1 expression could be found (Figure 18). For example, expression of splice variants I and IV was particularly strong in the lungs consistent with the association between AAA1 and asthma.

In the *in vitro* translation experiment, the *Xenopus* elongation factor α (pTRI-Xef) DNA template used as a positive control resulted in the synthesis of a 50-kDa polypeptide as expected. However, neither of the two investigated constructs was translated into polypeptides strongly arguing that AAA1 functions as a non-coding RNA gene (Figure 21).

To further investigate the translation of the AAA1 gene, a polyclonal antibody against the constant region (YVRRNAGRQFSHC) of the gene product was produced in rabbits. AAA1 peptide synthesis and antibody production were purchased from Sigma-Genosys Ltd (London Road, Pampisford, Cambridge). To test the specificity of the antibodies, Glutathione S-transferase (GST)-fusion proteins for AAA1 were produced with the pGEX 4T-3 GST fusion expression vector (Amersham Biosciences) according to the manufacturer's instructions. (Figure 21). The antibody displays high affinity against the recombinant AAA1 protein produced in bacterial lysate with no cross-reactivity between the GST construct alone. In spite of that, the antibody did not reveal any reactivity either in Western blots (spleen, skeletal muscle, uterine muscle, colon muscle, colon epithelium, kidney, testis and prostate) or in immunohistochemistry (bronchial tissue, HepG2 cell line) (data not shown).

In addition, formalin fixed, paraffin-embedded specimens of normal adult human bronchus tissue and HepG2-cells (positive for AAA1 RT-PCR) were used for immunohistochemistry. Immunohistochemical analyses were performed using rabbit AAA1 antibody and the ABC method (Vectastain Elite ABC kit, Vector Laboratories, Burlingame, California). Finally, AAA1 does not appear to have a mouse counterpart. Concluding from our data, AAA1 may be a non-coding RNA gene and is unlikely to encode a functional protein.

C. The Predicted Structure of the AAA1 Peptides

The cDNA sequences of twelve AAA1 splice variants are shown in Fig. 20. All predicted AAA1 proteins are small peptides (size from 34 aa to 74 aa) that do not show significant

identity to any known modular structures or motifs. All isoforms contain the same core sequence (AYVRRNAGRQFSHCNLHAHQFLVRRKQ) flanked by alternative amino- and carboxyterminal tails (Figure 15).

2. Characterization Of AAA1 as a Genetic Regulator of Asthma Related Traits

A. Association Analysis

AST1 covered the exons 3 to 10 of AAA1 (Figure 13). For the association analysis we chose one exonic SNP (SNP 517278) and four additional SNPs near exon-intron boundaries of AAA1 (Table 12). All the polymorphisms were identified previously (Table 3). The analysis for high total serum IgE level was done in the same data set as previously (304 disease associated and 220 control chromosomes, Table 2 and 8). Best haplotype associations of AAA1 are shown in Table 12. The best observe associations reached the χ^2 values of 8.9-13.6 and the permutation test showed statistically biased haplotype distribution for high IgE. Based on 10,000 simulations, the best marker-wise P value ≤ 0.0001 was observed for SNP_517278 in exon 10. Corrected P value for testing multiple markers simultaneously was 0.0002.

B. Variable alternative splicing for AAA1 depending on genotype

To study further whether AST1 effects on the expression of AAA1 we studied lymphoblast cell lines from the asthma patients who were homozygous, heterozygous, or non-carriers of AST1. Using RT-PCR with the primer pair SCF10 and ASKAR (Table 9) significant differences were found between patients and the differences were dependent on genotype (Figure 19). Only the non-carrier of AST1 processes normal amount of the exon 6-10b transcript, whereas the homozygote and heterozygotes show either an absent transcript or smaller splice variants. Differences in expression patterns suggest that AST1 can effect splicing of the gene and thereby, increase the risk of asthma related diseases among AST1 carriers compared to that among AST1 non-carriers.

REFERENCES

- Dantzker, D.R. *et al.* 1987. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease (COPD) and asthma. *Am. Rev. Respir. Dis.* **136**: 225-243.
- Becker, K.G. *et al.* 1998. Clustering of non-major histocompatibility complex susceptibility candidate loci in human autoimmune diseases. *Proc. Natl. Acad. Sci. U.S.A.* **95**: 9979-84.
- Daniels, S.E. *et al.* 1996. A genome-wide search for quantitative trait loci underlying asthma. *Nature* **383**: 247-50.
- Dizier, M.H. *et al.* 2000. Genome screen for asthma and related phenotypes in the French EGEA study. *Am. J. Respir. Crit. Care Med.* **162**: 1812-8.
- Jacob, H.J. *et al.* 1992. Genetic dissection of autoimmune type I diabetes in the BB rat. *Nat. Genet.* **2**: 56-60.
- Laitinen, T., *et al.* 2001. A susceptibility locus for asthma-related traits on chromosome 7 revealed by genome-wide scan in a founder population. *Nat. Genet.* **28**: 87-91.
- Leaves, N.I. *et al.* 2002. A detailed genetic map of the chromosome 7 bronchial hyper-responsiveness locus. *Eur. J. Hum. Genet.* **10**: 177-82.
- Malerba, G. *et al.* 2000. Linkage studies to asthma and atopy phenotypes on chromosomes 7, 12, and 19 in the Italian population. *Am. J. Hum. Genet.* **67**: 330.
- Mathias, R.A. *et al.* 2001. Genome-wide linkage analyses of total serum IgE using variance components analysis in asthmatic families. *Genet. Epidemiol.* **20**: 340-55.
- Ober, C. *et al.* 2000. A second-generation genomewide screen for asthma-susceptibility alleles in a founder population. *Am. J. Hum. Genet.* **67**: 1154-62.
- Polvi A *et al.* 2002. Physical map of an asthma susceptibility locus in 7p15-p14 and an association study of *TCRG*. *Eur. J. Hum. Genet.* **10**, 658-665 (2002).

Remmers, E.F. *et al.* 1996. A genome scan localizes five non-MHC loci controlling collagen-induced arthritis in rats. *Nat. Genet.* **14**: 82-5.

Satsangi, J. *et al.* 1996. Two stage genome-wide search in inflammatory bowel disease provides evidence for susceptibility loci on chromosomes 3, 7 and 12. *Nat. Genet.* **14**: 199-202.

Sawcer, S., *et al.* 1996. A genome screen in multiple sclerosis reveals susceptibility loci on chromosome 6p21 and 17q22. *Nat. Genet.* **13**: 464-8.

Sevon, P. *et al.* 2001a. TreeDT: Gene mapping by tree disequilibrium test. In KDD-2001. Proceedings of the Seventh ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, pp 365-370. Editors: Provost, F. and Srikant, R. Publisher ACM press, San Francisco, CA USA
(www.acm.org/sigkdd/kdd2001)

Sevon, P. *et al.* 2001b. TreeDT: Gene mapping by tree disequilibrium test (extended version). In the publication series Report C-2001, Nr 32:6-7 published by Department of Computer Science, University of Helsinki

Toivonen, H.T. *et al.* 2000. Data mining applied to linkage disequilibrium mapping. *Am. J. Hum. Genet.* **67**: 133-45.

Wjst, M., *et al.* 1999. A genome-wide search for linkage to asthma. German Asthma Genetics Group. *Genomics* **58**: 1-8.

Xu, J. *et al.* 2001. Genomewide Screen and Identification of Gene-Gene Interactions for Asthma-Susceptibility Loci in Three U.S. Populations: Collaborative Study on the Genetics of Asthma. *Am. J. Hum. Genet.* **68**: 1437-1446.

Xu J *et al* (2000). Major genes regulating total serum immunoglobulin E levels in families with asthma. *Am J Hum Genet.* **67**:1163-73.

- Yokouchi, Y. *et al.* 2000. Significant evidence for linkage of mite-sensitive childhood asthma to chromosome 5q31-q33 near the interleukin 12 B locus by a genome-wide search in Japanese families. *Genomics* 66: 152-60.
- Daly, M.J. *et al.* 2001. High-resolution haplotype structure in the human genome. *Nat Genet.* 29: 229-32.
- Dantzker, D.R. *et al.* 1987. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease (COPD) and asthma. *Am. Rev. Respir. Dis.* 136: 225-243.
- Johnson, E.N. *et al.* 2002. Heterotrimeric G protein signaling: role in asthma and allergic inflammation. *J Allergy Clin Immunol.* 109: 592-602.
- Kauppi, P. *et al.* 1998. Verification of self-reported asthma and allergy in subjects and in their family members volunteering for gene mapping studies *Resp. Med.* 92: 1281-1288.
- Laitinen, T. *et al.* 1997. Genetic control of serum IgE levels and asthma: Linkage and linkage disequilibrium studies in an isolated population. *Hum. Molec. Genetics* 6: 2069-2076.
- Laitinen, T. *et al.* 2001. A susceptibility locus for asthma-related traits on chromosome 7 revealed by genome-wide scan in a founder population. *Nat. Genet.* 28: 87-91.
- Michel U 2002. Non-coding ribonucleic acids--a class of their own? *Int. Rev. Cytol.* 218: 143-219.
- Numata *et al.* 2003. Identification of Putative Noncoding RNAs Among the RIKEN Mouse Full-Length cDNA Collection. *Genome Res.* 13: 1301-06.
- Palczewski, K. *et al.* 2000. Crystal structure of rhodopsin: A G protein-coupled receptor. *Science* 289: 739-45.
- Park, Y. *et al.* 2002. Identification of G protein-coupled receptors for Drosophila PRXamide peptides, CCAP, corazonin, and AKH supports a theory of ligand-receptor coevolution. *PNAS* 88: 11423-28.

Rana, B.K. *et al.* 2001. Genetic variations and polymorphisms of G protein-coupled receptors: functional and therapeutic implications. *Annu. Rev. Pharmacol. Toxicol.* 41: 593-624.

Although the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be clear to one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention. The above examples are provided to illustrate the invention, but not to limit its scope; other variants of the invention will be readily apparent to those of ordinary skill in the art and are encompassed by the claims of the invention. The scope of the invention should, therefore, be determined not with reference to the above description, but instead should be determined with reference to the appended claims along with their full scope of equivalents. Any embodiment, feature, step or element described above can be used in combination with any other unless otherwise apparent from the context. All publications, references, and patent documents cited in this application are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication or patent document were so individually denoted.

Table 1. Primers that were used in SNP genotyping with SBE method (A); primers, restriction enzymes, and the length of the digested PCR product of corresponding alleles used in SNP genotyping with altered digestion (B); primer pairs and the length of the corresponding PCR product used in genotyping of the small insertions and deletions (C).

Table 1A.

SNP ID	Nucleotide	Type ¹	Fw primer	Rev primer	PE PRIMER
dbSNP: 323926	NT 000380.3	C/T	CCCCCTGCTCTCTCTCTT	GGGATTATCCATTTCCTCCA	ATGCCAGCATCATCTTCCTG
dbSNP: 323906	509240	C/G	GATCTTCCATTTCACCAAGCA	GATGGATATTAGCTCCCCCTTA	CACACTAGCAAAAGTCAGGCAAT
dbSNP: 323917	515224	C/G	TGAGTTGATGAATCTGCTGGA	GGCAATGCCAATTAAAGAGGA	CACGAGCCCTGGATTACCTAC ²
dbSNP: 143266	515632	A/G	CAATGGATCACTGCAGCCTA	TTCCACCCCATTAGTTTTC	AGAACAAAGCTATGCCAGAGACTG ²
dbSNP: 324377	529556	A/C	AACACAGCCCTCCAGACACT	ATACCTGGCCAGCGGTTTA	TGGAAAAATTGAACATTATTGGG
dbSNP: 324373	531632	A/G	CTATTGGGCCCTGGAGACTT	ATGCAGCTTGCAATTGTTAC	GGTCTCTGGGACCCCTTC ²
dbSNP: 182718	543580	C/T	AGGGATCATGTCTCCAGCAC	TTAAATGCCCACTAGCACCA	GAGCAGAGATCAGAAACAGGCT ²
dbSNP: 324384	555608	A/G	TGTGTCTCCTTCTTGTGTTTATA	TGAAAGCAGGGCTACCATTT	GCAGGAACAGAAACGAAATACC ²
dbSNP: 324396	563704	C/T	CACAGGGAGGAAAGGAACAG	GATGGGATTGAGGAGCTTGA	ACAGTGGATGGTTTCTTGGC
dbSNP: 324981	591694	A/T	GGCCATCTGATAAAGCAGGA	TGATGCATAGGAATGCAAGG	AGTCTCCAGTGAATCGCCAA ²
dbSNP: 325456	617392	C/T	GATGAAAGGGAACCATTCCT	TGCTTTTCCTGTGTCCATTG	ATGCCCTAATTCTAAACCAATGA
dbSNP: 62911	638799	C/T	TGACATGTTCCCTGCATTG	AGAGGGCTTTGTGCTCTCTG	CACCCCTCCCTGGTGACCT ²

¹ Polymorphism detected in assays employing primers specified in this table

² extension primers (Pe) designed in reverse orientation

Table 1B. Primer, restriction enzymes, and sizes of the corresponding alleles after digestion used in SNP genotyping of AST1.									
Marker	Forward primer	Reverse primer	Enzyme	Product size*	Allele 1	Product size*	Allele 2	Note	
SNP_490331	gcctgccttagagacatcag	tggctgccttcttcattcttg	HpyCH4V	180+110	c	180+90+20	a	special combinations of fragments when heterozygous for the SNPs 490331 or 490390	
SNP_490390	gcctgccttagagacatcag	tggctgccttcttcattcttg	HpyCH4V	180+110	a	180+80+30	g		
SNP_490820	ccaattctcatccaaaggc	taaacctagggtgcgcacatttgcct cactggagatggc	Drall	330	t	250+80	g	plasmid tail with the Drall restriction site	
SNP_513659	aactgactaaactagggtgcacgtcg tatcctgacacagggatttaagc	tggatttgcatttgcgtctaag	MnlI	220	g	170+50	c	plasmid tail	
SNP_514743	ggaacatgggaagcaacacca	tctccctcagaaaaagccaga	SspI	420	g	210+210	a		
SNP_516094	ctgtcaataaacaattctcataattaaa	ggtggattacaggcatgag	FauI	400	t	210+180	c		
SNP_516174	gaagtggacttactgcatacaaga	gggtttcacctggttagc	SfiI	400	t	200+200	g		
SNP_516267	gctctagaaccccgctagcat	gcctcagcctctggagtagc	BsrBI	380	a	180+200	g		
SNP_520598	gcaaaagttagggggattc	accatgcctgcgcacatgtaa	NlaII	300	c	130+170	t		
SNP_521840	aggaacccttccacgacg	cccaagccctlgagaaaaat	BplI	360	a	180+180	g		
SNP_522363	tgacaccgcttttgaggta	tgtaacatcccccagggact	EcoNI	380	g	220+160	a		
SNP_525991	cttagtgggaaggctgtgg	aagccactaatatttggagtg	BsaWI	400	c	230+170	a		
SNP_528709	tgctcttttgacagccaatc	ggctgctttgatctgaggac	NcoI	360	t	180+180	g		
SNP_529142	ccataagcctgcctctattt	gtgggcacgtgagcagtagat	TaqI	410	a	240+170	g		
SNP_529820	caaatcctctgtctctctgct	taccagaggtctgagaaagg	BsaAI	400	t	220+180	c		
SNP_530177	ctttctcagcctctgggta	gtgagctgagttcgcgctat	StyI	380	t	230+150	c		
SNP_538567	ctcagaacactgctgcaga	gctctgggatgcgttaactt	EcoNI	380	g	220+160	a		
SNP_541908	ttgatctggccaaatggaac	ctaaacaaaaagggtgctgcaa	AflI	350	c	180+170	a		
SNP_546333	gcatgtgaatcacagcacttg	ctcccgagttcaagcaattc	HaeII	290+90	a	190+100+90	g		
SNP_563585	ccctacaagttacccttgcaca	cagaggttggatgggattg	HindII	375	c	180+185	t		
SNP_563930	tcaagctctcaatcccatc	gtggagggaagtgcctttca	StuI	330+40	g	150+180+40	a		
SNP_564782	aaaggcagccagtgcttgt	ggcagagagttccacagtga	MspI	400	a	170+230	g		
SNP_574953	tctcagcctgtaccctcta	aagcacaagcatcaggttca	AhdI	385	a	180+185	g		
SNP_585883	tcacacagggcatgattgga	gcagatggaggtagagctgtt	SfiI	370	g	180+190	c		
SNP_640764	aactgactaaactagggtgccacgtcg tcacatcatgaccatcgtgctctt	tagcacaatgcttgcctcat	SapI	340	t	290+50	c	plasmid tail	
SNP_647327	aggcaagtcacctgtctctc	gctgcagaaagaaccaaagg	BstEII	455	c	210+245	t		
SNP_662764	tgtctacctgttgccctg	gtctgtgcatctccaggt	MboI	455	a	240+215	g		
SNP_662803	catggagaagggaaggtcagg	ctagcacgtggcactgcctcta	DdeI	270	c	220+50	t		
SNP_663133	atgactgcatgacatgctta	tctgcaaacccagggtatct	SfaNI	400	t	140+260	c		

Table 2. HPM permutation test done using the following parameters: maximum pattern length 40 markers, one gap allowed for missing data and possible errors, and chi-square threshold for association 13.0. The observed scores (=number of qualified haplotype patterns spanning across the marker) and marker wise P values based on 10 000 permutations are shown. The critical region (=AST1) is highlighted.

Marker	HPM Score	Marker wise P value
NM4	0	1
NM5	0	1
D7S683	0	1
D7S656_2	0	1
NM2	0	1
MIT_MH26	0	1
NM8	0	1
NM9	0	1
NM11	0	1
G42099	0	1
SNP_490331	0	1
SNP490391	0	1
SNP_490820	0	1
SNP_506401	0	1
SNP_509240	0	1
NM50	0	1
SNP_513659	0	1
SNP_514743	0	1
SNP_515224	0	1
SNP_515632	0	1
SNP_516094	0	1
SNP_516174	0	1
SNP_516267	0	1
NM49_516568	0	1
NM51_517022	2	0.0139
G42102_518794	7	0.0105
NM53_519564	12	0.0091

Table 2. Continued

Marker	HPM Score	Marker wise P value
SNP_520598	15	0.0083
SNP_521640	20	0.0079
SNP_522363	23	0.0075
SNP_526991	23	0.0076
SNP_528709	30	0.007
SNP_529142	33	0.0066
SNP_529556	40	0.0062
SNP_529820	33	0.0061
SNP_530177	34	0.0059
SNP_531632	34	0.0054
D7S497_538277	40	0.0052
SNP_538567	38	0.0052
SNP_541906	36	0.0054
SNP_543580	35	0.0054
SNP_546333	35	0.0054
SNP_555608	35	0.0051
NM45_560804	35	0.0049
SNP_563585	35	0.0049
SNP_563704	28	0.0046
SNP_563930	23	0.005
SNP_564782	16	0.0063
SNP_574953	10	0.0071
SNP_585883	4	0.0093
SNP_591694	3	0.0098
SNP_617392	3	0.0085
SNP_638799	0	1
SNP_640764	0	1
G42097	0	1
SNP_647327	0	1
SNP_662764	0	1
SNP_662803	0	1
SNP_663133	0	1
NM13	0	1
NM46	0	1
D7S484	0	1

Genomic contig N1_000380 and corresponding alleles in haplotypes H1-H7

Type of polymorphism		Position in SEQ ID NO:1	Position in NT_000380	Polymorphism and flanking sequence		Alleles detected in haplotypes H1-H7							Location in GPR4 and AAA1 exons		Amino acid change
						H1	H2	H3	H4	H5	H6	H7			
SNP		1	509783	ACAATACAAATAGTTAAACCTAG[C>T]GGCAATTAATATGATGC	C	C	C	C	C	C	C	T			
SNP		93	509875	TGAGAGCATGAAAAAGAAAGGAC[A>G]AAAAATACAAAAAGA	A	A	A	A	A	A	A	G			
Insertion (NM50)		184-185	509886-67	GATAGAAACAACTTAAAGATAATAGTGGCCATAAATTTTCCCAACTCT	no ins							ins			
SNP		918	510700	TCCCCTGCCAATACAAAATTCGAG[G>A]ATGCTTAAGTCCCTTACATAAA	G	G	G	G	G	G	G	A			
SNP		983	510765	CTACATACACCCCTCTGTAT[A>T]CTTTAAGTCATCTCTAAATTAATT	A	A	A	A	A	A	A	T			
SNP		987	510769	CTACATACACCCCTCTGTAT[A>T]CTTTAAGTCATCTCTAAATTAATT	T	T	T	T	T	T	T	C			
deletion		1202-1208	510984-88	ATATTATATAAATCAATCTAAGT[TAAGACAGGAGAGAGAAAAATATTA	no del							no del			
SNP		1542	511324	TATCAAAAGTTTATAAATTTGCTGTGAT[C>T]TTGGAATCACTGTAGCTTATC	C	C	C	C	C	C	C	T			
SNP		1710	511492	CACCAATATATACCTATGATTTTGT[A>G]TACTTTTCTTATGTGTTTAT	A	A	A	A	A	A	A	G			
SNP		1818	511600	CAATGGCTTTAGGAATCAAGCAGG[C>T]AATGAATGTAACATGTAAA	C	C	C	C	C	C	C	T			
SNP		1927	511709	AAGACATTAATCAATTTTGT[C]>T]AACACTGTGCTGATCAACAAAT	A	A	A	A	A	A	A	T			
SNP		2254	512038	ATTTCTCTCCTAATGTGTT[C>A]TATTGTTTATTAGTATAAT	T	T	T	T	T	T	T	C			
SNP		2937	512719	CAACAACACAGAAATTTATTCCT[C]>A]CAGTCCTGTAGGCTGAAAGTTT	G	G	G	G	G	G	G	A/G			
SNP		3877	513859	TCACACAGGATTAAGCCT[G>C]AAGTTTTTTCAGAGAAGA	G	C	C	C	C	C	C	G			
SNP		4012	513794	TTACACCTTCAAGATGGCTATTC[C]>A]CTATAAATAATACTAA	C	A	C	C	C	C	C	A/G			
SNP		4631	514413	CAACAGCAATGCAAGAGAGAGATTT[C>G]TCGGAATCAGCTGCCTAAA	T	T	C	C	C	C	C	A			
SNP		4689	514471	CAAAAGAAATCAATGGCATATTC[C]>G]TCTCCCTGGGAGTTTATTA	C	C	C	C	C	C	C	G			
SNP		4981	514743	CTGTGGCACCCTCAATGAACATAAT[A>G]TTAAATGAATAT	A	G	A	G	A	G	A	G			
SNP		5442	515224	TATCTTATGTAAGAGAGTCCGGA[C>G]GTAGGTAATCCAGGGCTCGTGTG	C	C	C	C	C	C	C	G			
SNP		5634	515416	CAGGAAGGAAGAGGGAGGA[G>A]GACAGAGGGGCTCATATCAGCTCTTG	G	A	G	A	G	A	G	A			
SNP		5850	515632	AGAAATAGAAATTAAGGA[A>G]CAGTCTCTGGCATAG	A	G	A	G	A	G	A	C			
SNP		6312	516094	AATGTCGGGGACAAATGCTCTAGAA[C>T]CCGCTAGCATAGAC	C	T	C	T	C	T	C	G			
SNP		6392	516174	ACTGCAGTAGCTACA[G>T]AATCATAACTGGTTTTT	G	T	G	T	T	T	T	G			
SNP		6485	516267	AATTTGCTTATTATCAAAAG[G>A]AGCGCCAGCGCGGTGG	G	A	G	A	G	A	G	G			
SNP		6522	516304	TGGCTCATGCTGTATCCCA[C>G]CACTTTGGAGGCCAAGCGGG	C	G	C	G	A	G	C	C			
SNP		6646	516428	GGCGTGGTGGCGGCGG[A>G]TGGCAGCGCCTGTAGTCCAGC	A	G	A	G	A	G	A	A			
SNP		6739	516521	CGAGCTCGCAGTGCAGGATGCG[G>A]CCACTGCCCTCCAGCCTGGG	G	A	G	A	G	A	G	T			
SNP		6760	516542	ACTGCCCTCCAGCCTGGT[C>G]ACAGAGCAAGACTCC	T	C	T	C	T	C	T	T			
AAA-repeat (NM49)		6786-6817	516568-89	AGCAAGACTCCGCTCAAA[A]TAA[A]B[TA]A[7]GCAATTTTTTCACTTTAG	8							8			
deletion		6821	516603	ATAAATAAATAAATAAGCAATTTTTTTTTTCACTTTAGCCAGC	no del							no del			
SNP		7125	516907	TGAAGCCACTATTTTGG[C>A]ACATCTGTAATTTCTAGAA	C	C	C	C	C	C	C	C			
SNP		7229	517011	CCAAATTAATTAATTCATTTAC>T]CATCTTTAAGACTTACTTGCTGTTTA	C	T	C	T	C	T	C	T			
deletion (NM51)		7240-7243	517022-25	CCAAATTAATTAATTCATTTACCATCTTTAAGTACTTACTTGCTGTTTT	no del							no del			
SNP		7277	517059	ATGCAATAAAATG[C>G]GGAACATAAAGAGTAA	C	G	C	G	C	G	C	T			
SNP		7303	517085	AAGAGTAACCTCCGAT[G>T]AGTGTCAACAAGAAAGAAAGAA	G	T	G	T	G	T	G	C			no change (3'UTR)
SNP		7305	517087	AAGAGTAACCTCCGAT[G>T]AGTGTCAACAAGAAAGAAAGAA	G	T	G	T	G	T	G	C			no change (3'UTR)
deletion (NM52)		7306-7308	517088-90	AAGAGTAACCTCCGAT[G>T]AGTGTCAACAAGAAAGAAAGAA	no del							no del			no change (3'UTR)
deletion (NM52*241)		7334-7335	517116-17	CAACAAGAAAGAAAGAAAGAAACATATTTTTTGGCGCACATTCAATC	no del							no del			no change (3'UTR)
SNP		7486	517278	ACCTCCTATACATTACCTGAA[C>T]GCAGAGAAACTTGAGACGT	C	T	C	T	C	T	C	C			no change (3'UTR)
SNP		7550	517332	AGGTAAGGCGCAGGAGTGTGAGG[C>A]TCCAGGTCGCTGTGGAGTT	G	A	G	G	G	G	G	C			no change (3'UTR)
SNP		8490	518272	CTCTATATTGCTCTTTTATATA[C>T]ATATTTTTTGGCTGTTAACGTATG	C	T	C	C	T	C	C	C			no change (3'UTR)
T-repeat (G42102)		9012-9035	518794-817	TGTTGTGATGTATGTTGAT[C>T]12>[CT]10]CATACACACACATAAGCTTTT	12							12			
deletion		9199-9201	518981-83	TTTTGTAAGTGTGTTGGCATTATCTTCTTCAATGTGACCTTACITGAAA	no del							no del			
insertion		9355-9358	519137-38	TTGGGTAGTATTTTTTTTTTCAAGAGTCAITGGATAAGA	no ins							ins			
SNP		9649	519431	TCTATTTTGGTCTGCTGTTATG[T>G]GAGTTCTCTGAGCTGCCATA	T	G	T	G	T	G	T	T			
deletion (NM53)		9782-9785	519584-87	CTTCAGGAACCTCACTGCTGCTGCTGCTATGTTAGTAAATTTGGCCTTT	no del							no del			
SNP		10818	520598	AGAAATAGCTGGCTTCTGGCCAT[C>G]CAGAGGAAGATGAGAAATAGA	T	C	T	C	T	C	T	C			
SNP		11858	521640	TTCCCCAAACACATGGGCTG[A>G]CAGGTAAGATGGATACCTT	A	G	A	G	A	G	A	A			
SNP		12581	522363	GAATTTGGCGGCTCAATAC[G>C]TTTCTCAGGAGTCCCTGGGG	G	C	C	C	C	C	C	C			
SNP		16845	526627	TCTAGAAATTCCTCTTGTAGGCCATCAG[T>C]AATCTCACTGAAATTAGAC	T	C	T	C	T	C	T	T			
SNP		16893	526675	TATAAATAAATAATAA[T>C]ATTATGCATATTTTAAATAT	T	T	T	T	T	T	T	T			
SNP		16980	526762	CAGAAATGAAGGCCAAATCTTTAG[T>C]GGGAAGGCTGTGGAGCT	T	T	T	T	T	T	T	T			
SNP		17147	526929	TGGCAAGCTCTGCATAGTCC[G>A]TGACTGGGAGATGGGAA	G	A	T	G	A	C	T	G			

Polymorphism and flanking sequence			Alleles detected in haplotypes H1-H7							Location in GPR4 and AAA1 exons		Amino acid change	
Type of polymorphism	Position in SEQ ID NO:1	Position in NT_000380	H1	H2	H3	H4	H5	H6	H7				
SNP	17209	526991	TAGGCCATCCACGGCCTGTG>AICCGGATCCCCTCCAGAAATCTCAAGAAA							C	C		
SNP	17435	527217	GGGTCAAGAAATTTAAGTGAATACJA>GJGTAGGTGGCTCATCTTTTC							A	A		
SNP	18383	528165	AGAAACACACCCACAGCTCCAGGCTG>GJATATATCTACAGTAATGATGTG							G	G		
SNP	18927	528709	CATGTTCACAATATCCCATG>TJATGGGTCATGTGATGTG							G	G		
SNP	18978	528760	TGAATGTGAGTTGAAGGGACTJA>GJGTCTACTCCAGACTGAGACTGTGAAA							A	A		
SNP	19288	528050	GAGCAGCACCTGGGTG>CJTGAATAGCACCTGGGAAGC							G	G		
SNP	18272	528054	CATGAGGAGCAGCACCTGGGTGGTGAJA>TJAGCACCTGGGAAGCTGAACC							A	A		
SNP	19360	529142	ACTACCATGAGAAGAAAGTCTG>AJAGGGAAGATAAAAGTCAGAA							G	G		
SNP	19452	529234	CCTTGGGAAAGAGTGAAGAGGTG>AJGCTCCAGAGCTGAGTCCA							G	G		
SNP	18871	528453	CACTGCCATTATCTTCCAAAGAGIG>AJAATGGACAGACATATAATTACTA							G	G		
SNP	19712	528494	AGCCATAAATGATGGC>AJAATAACACAGCCCTCCAGA							C	C		
SNP	19774	528558	TTTTGAACATTATTGGGJC>AJAATCACTGCCTTCTGAC							C	C		
SNP	20038	528820	CTTAGGACCATGCTGTGATAT>CJGTGAGTCTCTCAATGAA							T	T		
SNP	20089	528871	TTGAATATTATGATATCACCAGGT>AJTTTTTAAACCGCTGGCCAGGTATCC							T	T		
SNP	20309	530091	CTGCTTTGCACCAAGACAGAGTIG>AJTCAGGGGCAGATTACAGGAGTCTTAC							G	G		
SNP	20395	530177	CGGAGGCACTGCTGCT>CJTGGAAATGCTTTAATAAACA							T	T		
SNP	20789	530571	CTCCCGATCTCAGGTGATCTGCCIG>TJCCCTCAGCCTCCAAAGTGTGGGATT							G	G		
SNP	21850	531632	TTTTGACCTTTCAGATCCCTC>TJGAAGGGTCCACAGAGACCC							C	C		
deletion (NM54)	22122-22123	531804-5	TCTTTGCTGAGCCTCTGTTGTTGTGAGTCTGGAACCC							no del	no del	del/no del	no del
SNP	22475	532257	TGTCTGCTGGCTGGTGAAGAGT>CJGTCCAGCGTTGGCAGGCAGC							T	T		AAA1 exon 9
SNP	22493	532275	GTGAGAAAGGTGCCAGCGTTGGCAGGCJA>GJGCTGAGAGAAGTGGGGGA							A	A		AAA1 exon 9
SNP	22715	532497	GTGCATGCATATGTGAGACCTG>AJTGACCTTCTCAACAGAGGGA							G	G		AAA1 exon 9
SNP	22869	532651	AGCTAGATTGGTCAGAGCTTTAGT>CJGAAGGAGCAGGTGACC							T	T		AAA1 exon 9
SNP	22934	532716	AAAAATGCAATAACACACAGTCCCTAGAJA>TJAATGATGACATGCTGCTCTA							A	A		AAA1 exon 9
SNP	24007	533789	TGTGTGATGTGGGGTTT>CJGGTGTGTGATGGGGAT							T	T		AAA1 exon 9
SNP	24264	534046	TCACATATCTGCTTCTAATCATCAAGTCTG>TJGAAAGAAATCTGGATACTAT							G	G		
SNP	24869	534651	GGGCTCACCATTTCAAGCACAGT>CJAATTAGAACGAGACCC							T	T		
SNP	26198	535980	CTGCAGATAATGACAGTTC>TJGTATGTCAGCATCAACATGGA							C	C	CT	
SNP	26356	538138	GTGAGGAAGGCCATTTTAGCTC>TJGAAAGTGAAGTGTCCCATTA							C	C		
SNP	26875	538457	AGCAGATTACTATTTCTGGGAAGATTCTJA>GJTTTGACCCCAATTACTTACA							A	A		
TAAA repeat (G42100)	26929-26968	538711-50	CTAAATAGCA(TAAA)10>(TAAA)11GATGAATGTTAATTATCT							10	11	10/11	
SNP	27404	537186	TTATCCATTTGGTTGAACCT>TJGAAACCCCATCTATCTTCCA							C	C		
SNP	28197	537979	GTGATATAAGAAAGTGCTTA>GJAJCATAGTATCTAGTTACAGTAGGTGACC							G	G		
CA-repeat (D7S497)	28495-28564	538277-346	CTTCTAAGTACC(CA)8GACT(CA)13>(CA)12(GA)(CA)8CT(CA)8GAGCTCTC							13	13	17	
SNP	28770	538552	AATTTACTATGTCCCATCAACTGTC>TJCCAAATCCTCTACCCG							C	T		
SNP	28785	538567	CCCAATCCTCTACCIG>AJGGATAAGATCACGCTATCTT							G	G		
SNP	28858	538640	TTGAGACATCTACAATTGAGGAT>CJACTGAAGGACAGACACTAGGTAATC							T	T		
SNP	28868	538648	TTGAGACATCTACAATTGAGGATGAAAG>GJAJACAGACACTAGGTAATC							T	T		
SNP	31224	541008	TTGGGAGCACCTCACACACATTAAGTIG>AJCCCTTTCTCCACTTCAACTT							G	G		
SNP	31910	541692	AGCTCAGAAGCTCCAGGTAGAGCA>AJACGCTCTTGCCTTAATCTTTAAAAA							G	A		
SNP	32124	541908	AATAACCCCTCTGATGGGCTTC>AJAGACCCATGATAGCTATA							C	C	A/C	
SNP	32185	541967	TTCTCACAAGAAACTGATGGGCATGAAGAT>CJTGCACTGACTGAAGTGTCTC							T	T		
SNP	32978	542758	AATTATCAATTTGGAATGAT>CJATGATTAAGAAATAAAACAGATTGTTT							T	T		
SNP	33350	543132	CCACCCACACACCTGGCTC>AJAATGTTTATTTTAGTAGCAA							C	C		
SNP	33788	543580	CTGAACAGTCTTCCCTCA>AJA GCCTGTTTCT GATCTCTGCT							G	G		
SNP	34362	544144	AGAGGAGAGGGCAGGAGAGATAA>GJTAATAAGTTAAAGTTTCCAAT							G	G		
SNP	34716	544498	ACAAGTAAGATACACACAAAJA>CJTGCAATTCACCAGTAAACC							A	C		
deletion	34909	544691	AATGAAATATTTAAAGTACCCCTJGGGTTACATTTTGGTACTT							no del	no del	no del	del
SNP	35559	545341	GTTTGTAGAAATTCACAACTGATAIC>TJTAAGGTCTATCATGTAAACAGSGCTC							C	C		
SNP	36551	546333	ATTTAAGAAATCCATGCA>AJGCCAGGTGACGTGGCTCA							G	G		
SNP	36909	546691	CAAGGCTTTGCAATAGCIC>AJTATCCTGCATAAATAAAGTTAGTT							C	A		
SNP	37327	547109	ATAGAAACTGGGAGGGGAA>GJTGACAGAGATGATATTGA							G	G		
SNP	37415	547197	AGAGGAGTCATAGTCATGGIG>AJAGTGACAGATCAGATTGGT							G	G		
SNP	37685	547467	GAGTGACATCTGGGTATCA>GJTTTTTAAAGGTCCCCAGAT							A	A		

Asn>Ser

AAA1 exon 8

Type of polymorphism	Position In SEQ ID NO:1	Position In NT_000380	Polymorphism and flanking sequence	Alleles detected in haplotypes H1-H7							Location in GPR2 and 'AAA1' exons		Amino acid change
				H1	H2	H3	H4	H5	H6	H7			
SNP	37931	547713	AAAAAACACACTAAGGTATAAGTTC[C>T]TCAAATCTAACCTGCCCTCTCCAG	C	T	C	C	C	T	T		AAA1 exon 4	no change (SUTR)
SNP	37859	547741	ATCTAACCTGCCCTCTCCAGGC[C>T]ACAGAGCTGGGAGTGGCAATCAGC	C	C	C	C	C	T	T			
deletion	38850-38852	548632-34	TTATCTGGAGCCCTCTCCCTC[CTC]GTGTTTGTGACCTTAATAAATG	no del	del	no del	no del	no del	no del	no del			
SNP	39314	549086	ACCTATTGAACAATAAATTGTAATA[G>A]TATAAAATCTCAACATGAATAAA	G	A	G	G	G	A	A			
SNP	39343	549125	ATAAAAAATCTCAACATGAATAAAG[G>T]CAGAAATAGAAAATTATATATGC	G	G	G	G	G	T	T			
SNP	39927	549709	CTGCTTTTCTCTCTTTGTACCCACAC[CT>C]GGGGAACAAAGCTGAAACAAA	T	C	T	C	C	C	C			
SNP	45828	555808	AGCTCACACTTGTAAAGTGGGAACATG[C>T]GTTATTTGTTTCTGTTCTGCG	C	T	C	T	T	C	T			
SNP	50197	559979	GCTACCTGGATCCAAGATGG[G>T]AAAGAGTGAGGAGTATGC	G	T	G	G	G	T	T			
SNP	50334	580118	AGGTGGATCAGAGGTCAAGAGATC[G>A]AGACCAATCTGCGCAACGTGG	G	A	G	G	G	A	A			
SNP	50493	580275	GTGGAGGTTGGAGTGAGCGGAGAGTGC[G>A]CCACTGCACCTCCAGCCTGG	G	G	G	G	G	A	A			
SNP	50632	580414	AGGATTTGGGCATGAATAGGAGCT[G>A]CAGGTTGGAGAACATCACAA	G	A	G	G	G	A	A			
SNP	50835	580617	CCTGCAGCTGGTCAGGGATGAGAGT[C>G]AGAGTTGACTTATCTGTGC	A	A	A	A	A	C	C			
SNP	50955	580737	GCTGACAGCTAAGTGCAGG[G>C]TGGAAATCAGAGATATCTTAATAGAT	G	C	G	G	G	A	G			
CA-repeat (NM45)	51022-51049	580804-31	AAGAACCAGGCATCTTAGA[CA]BTA[CA]7>[CA]8[CGTGCAGCAGCCACCGC	wt	wt	wt	wt	wt	(CA)8	(CA)8			
SNP	51217	580989	TTAACTTCAGTTTGAACAC[G>A]ACAATCTTATTTTATATACAACTAC	G	A	G	A	A	A	A			
SNP	51476	581258	CCTCTGCCAGCTCACAGCA[G>A]AAGGGGCTCTAATTCAGAGAAAAGGA	A	G	A	A	A	A	A			
SNP	51538	581318	CCAGTCATGACCTGTTATG[C>T]TTTCATATAGGCATCAGTTAGACCC	C	C	C	C	C	T	T			
SNP	51681	581643	GACCAGACAGAAAGCAGCTCAGGA[C>G]GGGCCAGAGCATGAGACTAG	C	C	C	C	C	G	G			
SNP	51884	581668	GGCCAGAGCATGAGACTAGG[G>T]GTGGTGGCAGCTGGGTTTAT	G	G	G	G	G	T	T			
SNP	51975	581757	TGCATGCCATACAAAC[G>C]GGAAATCTACATTACTGA	G	C	C	C	C	C	C			
Insertion	52288-52287	582068-89	TAGCATAGGCCATTGCATTTCCTTC[CCCCCAGTACCTGAATCTGTGA	no ins	no ins	no ins	ins	ins	no ins	no ins			
SNP	52573	582355	TGCAATGTTTCATGACCCAAACAAAGCT[G>A]AGCTCTGTGTGGCACTCTCA	G	A	G	G	G	G	G			
SNP	52776	582558	TTTCAAGGATAAAGGAAG[G>C]GGTAGAAGGGGAAGTGGAGAGGT	G	G	G	G	G	G	G			
SNP	53803	583585	GGTTCTTTTGGGAAGCT[C>T]GCTGCTGGGTGTTCCGA	C	C	C	C	T	C	C			
SNP	53922	583704	GTGGATGGTTTCTTGGCT[C>G]TGGGGCTCTCTGGC	T	C	C	C	C	C	C			
SNP	54148	583930	CAAGCATCTTGCA[G>A]GGCCTTGTCTGTGAGGGG	G	G	G	A	T	A	T			
SNP	54199	583981	ACACACAGTAGGCCAGTCTCTGCT[C>C]GCACCTGCTCAGTTAGGCTGTGTGG	T	T	G	T	C	C	C			
SNP	54641	584423	AGATCCCAAGTGGCAAC[G>C]ATGAGATTAGACCGG	G	G	T	G	T	C	C			
SNP	54751	584533	GCAGAACTTGCCAAAGGTT[C>G]AGGTGAAGCAGCTGCA	T	T	T	C	C	C	C			
SNP	55000	584782	TTGTGCTCCTGTTCTTCCC[G>A]GAGAATCGGCGCTG	G	G	G	A	G	G	G			
SNP	55134	584916	AGTGGGGAGCATTTGGGAGAGCTCTTC[G>A]GGTCCCTGACTGGTTCTCT	G	A	C	G	T	C	C			
SNP	56863	586465	AATAAAGCCAGAAATA[C>T]GAGAGCTGCTTCCCCAGT	C	C	C	C	C	C	C			
SNP	56858	586838	CCAGGTGTGGTGGTGCACGC[C>T]GTAGTCCAGCTATTCAGGAGGCTGAG	C	T	C	C	C	C	C			
SNP	57790	587572	TGGTTGGAGAAGGGTCCCTTCTTA>C]ATTTAGTTTATTTTATTTTATTTG	A	A	C	A	A	C	A			
SNP	60559	570341	TATTTGATTGCACTTA[G>G]AGTGAATAATATTAATTAATTA	C	C	C	C	C	G	G			
SNP	60804	570386	CTATTTAATTACAATTGCA>T]TCAAGCTCCAATAAACAATCC	A	A	A	A	A	A	T			
SNP	61165	570947	AGCTTTCTCTGGCCATGC[G>A]ACTGTTCTGCTAGTCTGCTC	G	G	C	C	C	C	C			
SNP	64559	574341	CTGCATTCTTTGGGAAGCGGATG[C>T]TGAACAGAGTCAGGTATACAAAAT	C	C	C	C	C	T	T			
SNP	65171	574953	CATCCTTTAAAGAGAA>G]ACAAGTTGCTATGTC	A	G	A	G	G	T	G			
SNP	65857	575639	AATTTAATTGGTAACTACCT[C>T]GACATCAGGTGGGCCATTACGTTTTAT	T	G	G	T	T	T	T			
SNP	66184	575946	TCCCTTCAGGAAGCCAGAGAGTGCATCTG[C>T]AACACCACA	C	C	C	C	C	T	T			
SNP	575972	575972	ACCACAATTAGGATAAAGGA[C>T]CTTATATATACAGAAATTTCAATGTCT	C	C	C	C	C	T	T			
SNP	576308	576308	ACTGAGTATCTGCCAGGCA[C>T]TATGAAAAGATAAACTATAAACATGA	C	C	C	C	C	A	A			
SNP	66628	576684	AAAATAATAATCTCTGTATCCCAT[A>G]GTATAGTGAATAGGAAGAAATA	A	G	T	T	T	C	A			
SNP	66902	577639	TAAATAAGTTTTATTGGAT>A]TACAGCCAGGCCCATCC	T	T	T	T	T	A	A			
SNP	67857	577701	TCCACTCTACAATGGCAGAT>C]TTGAGTAGCTGTTA	T	T	T	T	T	A	T			
SNP	67919	577701	AATAAGGTAAATAAGGAGGATGC[G>A]GTGACTGAGTATTGAGATGTTCTGA	G	G	G	G	G	T	G			
SNP	72270	582052	CAGTGTGAAGGCAGTCGA[G>A]CAGTGCATCTCTATAGTATGG	G	G	G	G	G	A	A			
SNP	75115	584897	CAGCCCCCATTCACCTCCACAA[C>T]AGCAGGGGCGAGGCA	C	C	C	C	C	C	C			
SNP	76101	585883	ACAACAGCAGGGGCGAGGGCCACCTA[G>C]AGCATAAATAAT	G	C	G	C	C	C	C			
SNP	81812	591694	ACTGGTCAACATCTGACAGATAATA>T]TTGGCGATTCACTGG	A	T	T	T	T	C	G			
SNP	82203	591985	GAGCCACAAATGAGCAATGCTAAGGAC[A>G]TAAAGCATATTTCAAAATG	A	G	A	T	A	G	G			
SNP	82332	592114	CTTTCACCTCTCAAAATGAACCTCAACG[C>T]TGGCTACTGCTCTTTTGTAAA	C	C	C	T	C	C	C			
SNP	82922	592704	TTGTATATATATTCAATAAGTAAATTT>A]TAAATTTTTTCATTGGTTTAT	T	T	T	T	T	T	T			
SNP													Asn>Ile

Position in Polymorphism and flanking sequence				Alleles detected in haplotypes H1-H7							Location in GPRA and AAA1 exons		Amino acid change
Type of polymorphism	Position in SEQ ID NO:1	Position in NT_000380		H1	H2	H3	H4	H5	H6	H7			
SNP	83552	593334	AAGTGGCAACATATTTATGAAC TAGT[C>T]TCAAAGGTCATACACCATCAGTT	C	C	C	C	C	T	T			
SNP	85227	595009	GCTCACCACATGTTTTTAACAT>C/CACAGCCAGCACCACATAGTAA	T	C	T	C	C/T	C	C			
SNP	85271	595053	AAGCTAGTTAACACTCAAGCCCA[G>A]GTGTTAACTCTGCCATGTGTC	G	G	G	G	G	A	A			
SNP	107610	617392	CCTAATCTAAACCAATGAT>C/ATTAAAGAAAGGAAACTA	T	C	C	C	C/T	C	C			
SNP	110989	620771	AATGCCCTCTTTAATTTTACTTT>C/CTTGATATACCTTTGGCCATCCCTCT	T	T	T	T	T	C	C			
SNP	111012	620794	CTTTCTTGATATACCTTTGGCCATCC[C>T]TCTATTTCCACACTTAGATATCT	C	T	C	C	C	C	C			
SNP	112030	621812	TATTCATGCTCTTCCCTCTCTT[G>C]AATTCCTAATTTAATTC AATTC TC	G	G	G	G	G	C	C			
SNP	112037	621819	TCTCCCTCTCTTGAATTCCTT>G/AAATTTAATTC AATTC TC	T	G	T	T	T	T	T			
SNP	112283	622065	TTCCTTTAGGTAACTTTTAA[A>T]TTTTTTACCCCTGCCCTTAA	A	T	T	T	T	T	T			
SNP	112726	622508	CCAAGAGATAATTTTTCTTT>ATTAGTTTTTCAGAGGAGCCCTTC	T	T	T	A	A	T	T			
SNP	112859	622641	GTTTGTGTACAGATTATTTA>C/ATCACCCAGGATTTTTTTT	A	C	A	C	C	C	C			
SNP	113428	623210	TAATCATTCTCAGAACCTCT[C>G]CCTATGGTGGCTTTCACAACATTAGTG	C	C	C	C	C	G	G			
SNP	113845	623427	TCTACTCCCTCTGGAATTTGA[A>T]CCTAATGTTACAAAGTGG	A	T	A	T	A/T	T	T			
SNP	113944	623726	AGTGGATGAAAAGGGGAGCT>G/CTCTGATGGTGCAGTGCTG	T	T	T	G	G	T	T			
SNP	114945	624727	GGAAATACCTGTTCTCATTTA[A>G]GACCACTGGTTTCCAGGCTT	A	A	A	G	A/G	A	A			
SNP	115192	624974	CATTTCTGCTTTAGGTTGTGCTGCT[C>G]TACGCCCTCTACCTACGTCCTGGT	C	G	C	C	C	C	C			
SNP	115828	625410	CATGCATACCTGTCCAAAGAAAT>C/GAACTCTCCACACTAATCTCTCATTT	T	T	T	T	T	C	C			
SNP	116032	625814	GGCCCTGGGAGCAGTAAT[G>A]CCCTAGAGGAGGGTGAGTA	G	G	G	A/G	G	G	G			
SNP	116464	626246	CTTATTGAAATATAAATATG[G>A]CAGCCCTTTGATATGACTTC	G	G	G	A	A/G	G	G			
SNP	116515	626297	TCTGGAGAAACACCCACAC[G>A]TGTGCATGTATACATATGTG	G	G	G	G	G	A	A			
SNP	116926	626708	TAGAGCCATGGGCAGTAAT>C/GCATCAACATGAACATATC	T	T	T	T	T	C	C			
SNP	117276	627058	ATTTATTTTGATCATGCTTGT>A/TATGCCCTACTGTACAGCATAAGACATATG	T	T	T	T	T	A	A			
SNP	123667	633449	ACATTAAAGTGGTATGCATT[G>A]ATATGTATAGCTTTTTATAT	G	G	G	A	A	G	G			
SNP	123770	633552	TGTTTCTCTGGGTTCTCTCGGT[C[A>G]TTTGGTGTGCTGTTTGTGAAGGG	A	G	A	A	A	A	A			
SNP	123788	633570	GGTCATTTGGTGTGCTGTTT[G>C]TGAAGGGTTGAGACCTGTACCAAGTTT	G	C	G	G	G	G	G			
SNP	128017	638799	CAGAGAAAGAGGCCCTCCCTG[A>G]AGGTCACCCAGGGAGGGTGAAGG	A	A	A	G	nd	G	nd			

no sequencing/genotyping result is available for the site

del=deletion

ins=insertion

nd no sequencing/genotyping result is available for the site

del=deletion

ins=insertion

GPRA exon 4

no change

Table 4. Primers used in the cloning of GPRA.

Primer	Sequence	Exon	Splice variant
JEGE1F1	tctgtgctcctcggtcagcag	5' UTR	-
JEGE1F2	aagctggactccctcactcagc	5' UTR	-
JEGE1F3	agcaaggacagtgaggctcaacc	5' UTR	-
JEGE9aR1	ctggcatgaataactggggagttc	3' UTR	A
JEGE9aR2	tttgtcttgatcatctcccaggta	3' UTR	A
JEGE9bR1	tatagccctccctgggtgaatctga	3' UTR	B
JEGE9bR2	gtgccctggtaagcagtgagaagt	3' UTR	B
JEGE5F1	tgagcaattgataactctgtgggtcctc	E2a	-
JEGExR2	AAATAAGCTGTGGCATCCTCATCCAG	3' UTR	C
JEGExR1	TGGCATCCTCATCCAGGGATATTGC	3' UTR	C
VAUE3R1	GGAAGGCCACGATGGTCATGTATGG	E5	-
Vau8R1	ATGACGAGGGTGGGTGAAGTTGG	3' UTR	A
Vau8R2	GAATGGTGGGGAAGGAAGGCGTTT	3' UTR	A
Vau1000R1	ggccatcctgctgtgacccattt	3' UTR	B
AS8FX1	ATGAGATGCAGATTCTGTCCAAG	E9a	A
AS8FX2	TGCACAAGACAAATGTTCTAATGA	3' UTR	A
AS8F5.1	CAGCTATAACCGAGGACTCATCTC	E7	-
AS8F7.1	TGTTGGAGTCCATACTTCTGTT	E8	-
VAUE3R2	AAAAGACAGGCTCCAGGCGATCACA	E5	-
A2E2F	gattctcccagtggtgcacgaa	E1	-
A3E3R	TGATGGCCAGCTGAGTCACAAAGAAGG	E2a	-

Table5. Primer pairs used in re-sequencing of the exons and exon/intron boundaries of *GPRA*

Primer	Forward primer	Reverse primer
<i>GPRA</i> ex1	actcagctgcaggagcaag	tgacactcttaagttccagcagtc
<i>GPRA</i> ex2a	aggaggaagaaatccagcct	tgaccgatgcgttacatttt
<i>GPRA</i> ex2b.1	gccatacattgtagtaacctgaaa	cctcatccagggataatttgc
<i>GPRA</i> ex2b.2	ttcctaccaacaagaactccaa	cgatgatgaattagaacatacaacttt
<i>GPRA</i> ex3	taagtcaaagaactcctaccttgc	agcaaaggaaatacattaaaaatcaaa
<i>GPRA</i> ex4	ctgccctctttcaccagta	agccaccaccttccttagt
<i>GPRA</i> ex5	gcttctgttcaagcttcccttt	gtgtggtcctgtcctgacg
<i>GPRA</i> ex6	tggatcctcatggtcactttc	tctctgctggcatagcttga
<i>GPRA</i> ex7	ttgagagagtctgagcattcca	ccaaattattcaacccatagcc
<i>GPRA</i> ex8	tgacatcaatgctccaaacaa	tgtcatgattaaggcggtttc
<i>GPRA</i> ex9a	ttaacatgtctacttgccctttca	tctgcaaaccgaggctatct
<i>GPRA</i> ex9b	gcagagctgtcacccaaaat	agcctgggcaacaagagtaa

Table 6A. Primers used in *GPR4* SNP genotyping.

SNP ID	Nucleotide	Type	Fw primer	Rev primer	Pe primer
dbSNP: 324981	NT 000380.3 591694	A/T	GGCCATCTGATAAAGCAGGA	TGATGCATAGGAATGCAAGG	AGTCTCCAGTGAATCGCCAA

Table 6B. Primer pairs, restriction enzymes, and corresponding allele sizes used in genotyping of GPRA exonic SNPs.

Table 6B. Primer pairs, restriction enzymes, and corresponding allele sizes used in genotyping of GPRA exonic SNPs.									
Six digit number in the name of the SNP		shows its position in the genomic contig NT_000380.							
Marker	Forward primer	Reverse primer	Enzyme	Product size*	Allele 1	Product size*	Allele 2		
SNP_640764	aactgactaaactaggtgccacgtcgtccatcatcaccatcgtgctctt	tagcacaaatgccctgccctat	SapI	340	t	290+50	c		
SNP_662764	tgctacacctgtggcctgtg	gctctgtcatctcccaggt	MboII	455	a	240+215	g		
SNP_662803	catggagaaggaaggtcagg	ctagcacitggcacitgcccta	DdeI	270	c	220+50	t		
SNP_663133	atgacitcatgcacitgctta	tctgcaaacccgaggctatct	SfaNI	400	t	140+260	c		
* product sizes are approximations									

Table 7. SNPs found in the exons of six splice variants of *GPRA*.

Exon	Position of the SNPs						Sequence around the polymorphism	Amino acid change in different variants
	seq1	seq2, A variant	seq4, B long variant	seq6, B short variant	seq8, C variant	seq10, D variant	seq12, E variant	seq14, F variant
ex2b	-	-	-	-	585 -	-	-	-
ex2b	-	-	-	-	655 -	-	-	-
ex2b	-	-	-	-	681 -	-	-	-
ex3	81912	448	448	415	-	-	448	-
ex4	115192	524	524	491	-	420	-	-
ex5	-	776	776	743	-	672	682	578
ex6	-	-	-	-	-	-	-	-
ex9a	-	851	851	818	-	-	-	-
ex9a	-	1159	-	-	-	-	-	-
ex9a	-	1199	-	-	-	-	-	-
ex9a	-	1529	-	-	-	-	-	-
ex9b	-	-	1206	1173	-	-	-	-
ex9b	-	-	1225	1192	-	-	-	-
ex9b	-	-	1330	1297	-	-	-	-
ex9b	-	-	1338	1305	-	-	-	-

* at these sites contig NT_00380 contains an allele that associates with high serum IgE level and /or asthma.

Table 8. Best haplotype associations of *GPRA* for high serum IgE level in the data set of 304 disease associated and 220 control chromosomes when one gap (marked with an asterisks) was allowed in the haplotype patterns.

High IgE	Control	Conf	χ^2	<i>GPRA</i> markers				
				SNP591694	SNP640763	SNP662763	SNP662803	SNP663133
				Exon 3	Exon 5	Exon 9A	Exon 9A	Exon 9A
51	16	0.761	10.3	-	C	-	-	-
44	14	0.759	8.5	T	C	-	-	-
40	12	0.769	8.5	-	C	*	T	-
39	12	0.765	7.9	-	C	*	T	C
39	12	0.765	7.9	-	C	G	-	-
39	12	0.765	7.9	-	C	G	T	-
38	12	0.76	7.3	-	C	G	*	C
38	12	0.76	7.3	-	C	G	T	C
35	11	0.761	6.8	T	C	*	T	-
34	11	0.756	6.2	T	C	*	T	C
34	11	0.756	6.2	T	C	G	-	-
34	11	0.756	6.2	T	C	G	T	-
33	11	0.75	5.7	T	C	G	*	C
33	11	0.75	5.7	T	C	G	T	C

High IgE, number of disease-associated haplotypes with a specific pattern; Control, number of control haplotypes with a specific pattern; Confidence, percentage of haplotypes with the specific pattern that is associate with disease, chi-square value for disease association of the specific haplotype pattern.

Table 9. Primers used for cloning of full length cDNAs for AAA1.

Primer	Exon	Sequence
IF	1	5' AGAATGAGTCTCTGATGACTTT 3'
IIF	2	5' ACTTGCTGTTTCATAGAATTGCAA 3'
SF9	4	5' TGACTTCTCCCCAGATTTTTGTAT 3'
exIF	5	5' ACACATACAAAGTGCCTACCACAT 3'
SR13	9	5' TTGAAACTGTATTTCCCATATTGC 3'
ASKAF/R	10b	5' AAATGCAATAAAAAATGCGGAACTA 3'
XR	11	5' GAGTCATTAGTCCAGAGAACAT 3'
XIR	12	5' CTGCTTGGAACAGTGTATATC 3'
XIIR	14	5' TGGTCTACGTAGAATTCAGAGTA 3'
XVR	16	5' CATGTGTTAATTGTGTCTTCACT 3'
SR4	19	5' GGGTGTCATTTACACGAACAATAA 3'
SF10	6	5' CAGTTCAGTCACTGCAATCTTCAT 3'

Table10. Primers used in SNP genotyping with SBE method.

SNP	forward	reverse	extension
517278	TTATGAGCTAAAGTGCCCAATTAA	AGGAAGGTGGCAGTGAACTC	CCCTCCTATACATTACCTGAA
549709	TCCCGTCTCCTCTAGTCTTC	TGAGATGCTGTCTCTAAATAATAGA	TTCTCTCTTTGTACCCACAC
570341	TGTTGGTTCGATGAGAGCAT	AAGTGGGGAACAACACTGG	GTACCTATATTTGTATTGCACTTA

Table 11. Exon-intron structures and splice junction sites of AAA1. Gray area shows the exons and introns located in AST1.

Exon	Size (bp)	site in NT_00380	5' splice donor	Intron size (kb)	5' splice acceptor
1	84	684692 - 684776	TACTGGAAAGgtatgattgt	37.0	gattttacagGCTGATGGTG
2	189	647354 - 647543	CTAAGACCTGgtaagttat	66.0	ctactgtgagGACACAGAGA
3	99	581535 - 581634	AGGTCCTGCTGgtaaggaaat	7.0	ctatttcagAGTTCCTCAC
4	80	574305 - 574385	GAAGCAGGAGgtaaggagg	2.8	gottfoacagGAGTGACTTC
5	199	571268 - 571467	CATAGACCTGgtgagtaact	29.3	ctotggggagGCTCTAGGACT
6	80	541930 - 542010	AAGAAACAGgtaogtag	5.3	ccatttacagCTTATGTGAG
7	112	536477 - 536589	CTCTGGTATTTgtaagtaatt	2.5	gthttgacagGAGCTCCAAT
8	144	533835 - 533979	AAATACCTGCAGtaagtaca	0.8	oattgtacagTGAACAAATA
9	947	532055 - 533002	TGTCCATGAAggtctggac	14.7	catcttctagAAGTGAGAAA
10a	120	517273 - 517393	CTGCGTTTCAGgtaagtata	60.0	caaaatccagGAAACAAGG
10b	351	517042 - 517393	TATTGGATTT		taaaattagagGTTIAGGAAA
11	108	456437 - 456545	AGCTGGGTTGgtaagactgg	26.4	tttttgcagTGGGATATAC
12	97	430100 - 430197	CCATATATGAgtgatcata	47.2	cccttcgcagGACTGGCAAC
13	150	382905 - 383055	TCTGCCATGTgtaagtcttc	1.3	gattctgaagGCCATGTTTT
14	121	381445 - 381566	GGGACATTTTgttttccatt	1.1	tgaatttttagGACGCTGGGC
15	71	380274 - 380345	ATGACCTGAGgtgggtatta	0.3	ctgtccacagCAATCCTTCT
16	90	379916 - 380006	CGTTGTTACTaaaaaagaaa	149.0	ttctcaacagAGCAAACTGC
17	94	230772 - 230866	TGAGCCTCAGgtaagaaccc	66.7	ttgttttagGCTTTTGTAT
18	118	168107 - 168224	TATTTTAAAGgtacatgtat	4.1	ttctcccagCTCTGTTTTC
19	426	163615 - 164041	TTTCTAATTAAAAataaaa		

Table 12. Best haplotype associations of AAA1 for high serum IgE level in the data set of 304 disease associated and 220 control chromosomes when one gap (marked with an asterisks) was allowed in the haplotype patterns. Only the markers from the coding region or close to intron-exon boundaries were included.

Markers of the AST1 locus							
High IgE	Control	X ²	SNP ₅₁₇₂₇₈ (C>T) Exon 10a / 10b	SNP ₅₃₈₅₆₇ (G>A) Intron 6	SNP ₅₄₉₇₀₉ (T>C) Intron 5	SNP ₅₇₀₃₄₁ (G>G) Intron 5	SNP ₅₇₄₉₅₃ (A>G) Intron 3
59/187 (24%)	18/179 (10%)	13.548	T	A	-	-	-
49/190 (21%)	14/177 (7.9%)	12.5479	T	A	*	G	-
49/190 (21%)	14/177 (7.9%)	12.5479	T	A	*	G	G
117/130 (47%)	55/181 (30%)	12.5318	T	-	-	-	-
53/189 (22%)	16/177 (9.0%)	12.2929	T	A	C	*	G
53/189 (22%)	16/177 (9.0%)	12.2929	T	A	C	-	-
47/180 (21%)	14/174 (8.0%)	12.2377	T	*	C	G	-
47/180 (21%)	14/174 (8.0%)	12.2377	T	*	C	G	G
47/190 (20%)	14/177 (7.9%)	11.4627	T	A	C	G	G
47/190 (20%)	14/177 (7.9%)	11.4627	T	A	C	G	-
49/190 (21%)	15/177 (8.4%)	11.3004	-	A	*	G	G
103/136 (43%)	47/174 (27%)	11.2636	T	*	C	-	-
50/187 (21%)	16/177 (9.0%)	10.9929	-	A	*	G	-
47/189 (20%)	15/177 (8.4%)	10.3762	-	A	C	G	-
47/190 (20%)	15/177 (8.4%)	10.2633	-	A	C	G	G
60/184 (25%)	23/179 (13%)	9.0248	-	A	-	-	-
53/189 (22%)	19/177 (11%)	8.957	-	A	C	*	G

High IgE, number of disease-associated haplotypes with a specific pattern, Control, number of control haplotypes with a specific pattern, chi-square value for disease association of the specific haplotype pattern

Table 13. Haplotype specific AST1 polymorphisms for diagnostic testing. These markers can be used for the haplotype identification of an individual without phase information.

Haplotype	Type of polymorphism	Position in SEQ ID NO:1	Position in NT_000380	Allele present in the specific haplotype	Allele present in other haplotypes
H1	SNP	4631	514413	T	C
H1	SNP	51975	561757	G	C
H1	SNP	53922	563704	T	C
H2	SNP	5634	515416	A	G
H2	SNP	6739	516521	A	G
H2	SNP	7550	517332	A	G
H2	deletion	9199-9201	518981-83	deletion of TCT	no deletion
H2	SNP	16980	526762	C	T
H2	SNP	17147	526929	A	G
H2	SNP	17435	527217	G	A
H2	SNP	19272	529054	T	A
H2	SNP	19452	529234	A	G
H2	SNP	20309	530091	A	G
H2	SNP	20789	530571	T	G
H2	SNP	24869	534651	C	T
H2	SNP	32976	542758	C	T
H2	SNP	34716	544498	C	A
H2	deletion	38850-52	548632-34	deletion of CTC	no deletion
H2	SNP	50955	560737	C	G
H2	SNP	51476	561258	G	A
H2	SNP	52573	562355	A	G
H2	SNP	55134	564916	A	G
H2	SNP	56856	566638	T	C
H2	SNP	65857	575639	G	T
H2	SNP	66526	576308	T	C
H2	SNP	66902	576684	G	A
H2	SNP	72270	582052	A	G
H2	SNP	111012	620794	T	C
H2	SNP	112037	621819	G	T
H2	SNP	115192	624974	G	C
H2	SNP	123770	633552	G	A
H2	SNP	123788	633570	C	G
H3	SNP	22869	532651	C	T
H4	deletion	7334-35	517116-17	deletion of AT	no deletion
H4	SNP	16893	526675	C	T
H4	SNP	17209	526991	A	C
H4	SNP	26198	535980	C/T*	C
H4	SNP	32124	541906	A/C*	C
H4	SNP	36551	546333	A	G
H4	SNP	116032	625814	A/G*	G
H6	SNP	54199	563981	C	T
H6	SNP	57790	567572	C	A
H6	SNP	64559	574341	T	C
H7	SNP	1	509783	T	C
H7	SNP	93	509875	G	A
H7	SNP	918	510700	A	G
H7	SNP	983	510765	T	A
H7	SNP	987	510769	C	T
H7	SNP	1542	511324	T	C
H7	SNP	1710	511492	G	A
H7	SNP	1818	511600	T	C
H7	SNP	1927	511709	T	A
H7	SNP	2254	512036	C	T
H7	SNP	4689	514471	G	C
H7	SNP	5442	515224	G	C
H7	SNP	18383	528165	A	G
H7	SNP	18978	528760	G	A
H7	SNP	19671	529453	A	G
H7	SNP	28866	538648	C	G
H7	SNP	52776	562558	C	G

* A rare polymorphism found only in H4 haplotype

Table 14. AST1 polymorphisms specific for the different haplotype combinations for diagnostic testing. These markers can be used for the identification of certain haplotype combinations of an individual without phase information.

Haplotype combination	Type of polymorphism	Position in SEQ ID NO:1	Position in NT_000380	Allele present in the specific haplotype combination	Allele present in other haplotypes
H4+H5	SNP	7125	516907	A	C
H4+H5	SNP	18927	528709	T	G
H4+H5	SNP	19268	529050	C	G
H4+H5	SNP	19712	529494	A	C
H4+H5	deletion	22122-23	531904-5	deletion of TG	no deletion
H4+H5	SNP	24007	533789	C	T
H4+H5	SNP	27404	537186	T	C
H4+H5	SNP	28785	538567	A	G
H4+H5	SNP	33350	543132	A	C
H4+H5	SNP	33798	543580	A	G
H4+H5	SNP	34362	544144	C	G
H4+H5	insertion	52286-87	562068-69	insertion of CC	no insertion
H4+H5	SNP	53803	563585	T	C
H4+H5	SNP	55000	564782	A	G
H4+H5	SNP	56683	566465	T	C
H4+H5	SNP	67919	577701	C	T
H4+H5	SNP	76101	585883	C	G
H4+H5	SNP	82332	592114	T	C
H4+H5	SNP	112726	622508	A	T
H4+H5	SNP	113944	623726	G	T
H4+H5	SNP	114945	624727	G	A
H4+H5	SNP	116464	626246	A	G
H4+H5	SNP	123667	633449	A	G
H2+H4+H5	SNP	4012	513794	A	C
H2+H4+H5	SNP	4961	514743	G	A
H2+H4+H5	SNP	5850	515632	G	A
H2+H4+H5	SNP	6312	516094	T	C
H2+H4+H5	SNP	6392	516174	T	G
H2+H4+H5	SNP	6485	518287	A	G
H2+H4+H5	SNP	6522	516304	G	C
H2+H4+H5	SNP	6646	516428	G	A
H2+H4+H5	SNP	6760	516542	C	T
H2+H4+H5	deletion	6821	516603	deletion of T	no deletion
H2+H4+H5	deletion	7240-43	517022-25	deletion of ACTT	no deletion
H2+H4+H5	SNP	7277	517059	G	C
H2+H4+H5	SNP	7303	517085	T	G
H2+H4+H5	SNP	7305	517087	C	G
H2+H4+H5	deletion	7306-8	517088-90	deletion of TGT	no deletion
H2+H4+H5	SNP	7496	517278	T	C
H2+H4+H5	SNP	8490	518272	T	C
H2+H4+H5	SNP	9849	519431	G	T
H2+H4+H5	deletion	9782-9785	519564-67	deletion of GTCT	no deletion
H2+H4+H5	SNP	11858	521640	G	A
H2+H4+H5+H7	SNP	3877	513659	C	G
H2+H4+H5+H7	SNP	7229	517011	T	C
H2+H4+H5+H7	SNP	10816	520598	C	T
H2+H4+H5+H7	SNP	12581	522363	C	G
H2+H4+H5+H7	SNP	19360	529142	A	G
H2+H4+H5+H7	SNP	19774	529556	A	C
H2+H4+H5+H7	SNP	20038	529820	C	T
H2+H4+H5+H7	SNP	20395	530177	C	T
H6+H7	SNP	22475	532257	C	T
H6+H7	SNP	22493	532275	G	A
H6+H7	SNP	22715	532497	A	G
H6+H7	SNP	22934	532716	T	A
H6+H7	SNP	24264	534046	T	G
H6+H7	SNP	26356	536138	T	C
H6+H7	SNP	26875	536457	G	A
H6+H7	SNP	28197	537979	A	G
H6+H7	SNP	28858	538640	C	T
H6+H7	SNP	31224	541006	A	G
H6+H7	SNP	32185	541967	C	T
H6+H7	deletion	34909	544691	deletion of T	no deletion

Table 14. (Continued)

Haplotype combination	Type of polymorphism	Position in SEQ ID NO:1	Position in NT_000380	Allele present in the specific haplotype combination	Allele present in other haplotypes
H6+H7	SNP	37327	547109	T	G
H6+H7	SNP	37415	547197	A	G
H6+H7	SNP	37685	547467	G	A
H6+H7	SNP	37959	547741	T	C
H6+H7	SNP	39343	549125	T	G
H6+H7	SNP	50493	560275	A	G
H6+H7	SNP	50835	560617	C	A
H6+H7	CA-repeat	51022-49	560804-31	(CA)8	(CA)6TA(CA)7
H6+H7	SNP	51536	561318	T	C
H6+H7	SNP	51884	561666	T	G
H6+H7	SNP	60604	570386	T	A
H6+H7	SNP	61165	570947	A	G
H6+H7	SNP	75115	584897	A	G
H6+H7	SNP	82922	592704	A	T
H6+H7	SNP	83552	593334	T	C
H6+H7	SNP	85271	595053	A	G
H6+H7	SNP	110989	620771	C	T
H6+H7	SNP	112030	621812	C	G
H6+H7	SNP	113428	623210	G	C
H6+H7	SNP	115628	625410	C	T
H6+H7	SNP	116515	626297	A	G
H6+H7	SNP	116926	626708	C	T
H6+H7	SNP	117276	627058	A	T
H2+H6+H7	SNP	28770	538552	T	C
H2+H6+H7	SNP	31910	541692	A	G
H2+H6+H7	SNP	37931	547713	T	C
H2+H6+H7	SNP	39314	549096	A	G
H2+H6+H7	SNP	50197	559979	T	G
H2+H6+H7	SNP	50334	560116	A	G
H2+H6+H7	SNP	50632	560414	A	G
H2+H6+H7	SNP	51217	560999	A	G
H2+H6+H7	SNP	51861	561643	G	C
H4+H5+H6+H7	SNP	54148	563930	A	G
H4+H5+H6+H7	SNP	54641	564423	C	G
H4+H5+H6+H7	SNP	54751	564533	C	T
H4+H5+H6+H7	SNP	60559	570341	G	C
H4+H5+H6+H7	SNP	66164	575946	T	C
H4+H5+H6+H7	SNP	67857	577639	A	T
H2+H7	SNP	16845	526627	C	T
H2+H7	SNP	20089	529871	A	T
H2+H4+H5+H6+H7	SNP	21850	531632	T	C
H2+H4+H5+H6+H7	SNP	36909	546691	A	C
H2+H4+H5+H6+H7	SNP	39927	549709	C	T
H2+H4+H5+H6+H7	SNP	45826	555608	T	C
H2+H4+H5+H6+H7	SNP	65171	574953	G	A
H2+H4+H5+H6+H7	SNP	112283	622065	T	A
H2+H4+H5+H6+H7	SNP	112859	622641	C	A

Table 15. The cellular location of transiently expressed GPRA A, B, B-short, C, D, E, and F variants.
Results are means at least from two independent experiments.

	GPRA A	GPRA B	GPRA B-short	GPRA C	GPRA D	GPRA E	GPRA F
cytoplasm	29%	48%	100%	100%	100%	100%	100%
Plasma membrane	71%	52%	0%	0%	0%	0%	0%

WHAT IS CLAIMED IS:

1. An isolated GPRA polypeptide comprising an amino acid sequence that has at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ. ID NOS: 3, 5, 7, 9, 11, 13 and 15 over the entire length of the selected SEQ ID NO. when compared using the BLASTP algorithm with a wordlength (W) of 3, and the BLOSUM62 scoring matrix.
2. The isolated GPRA polypeptide of claim 1 comprising an amino acid sequence selected from the group consisting of B-Long (SEQ ID NO: 5), B-Short (SEQ ID NO:7), C (SEQ ID NO:9), D (SEQ. ID NO:11), E (SEQ ID NO:13) and F (SEQ ID NO:15) provided that up to 34 amino acids can be substituted, deleted or inserted relative to the selected SEQ ID NO.
3. The isolated GPRA polypeptide of claim 1 comprising the amino acid sequence selected from the group consisting of B-Long (SEQ ID NO: 5), B-Short (SEQ ID NO:7), C (SEQ ID NO:9), D (SEQ. ID NO:11), E (SEQ ID NO:13) and F (SEQ ID NO:15)
4. An isolated GPRA polypeptide comprising at least 10 contiguous amino acids from amino acids 343-377 of B-long (SEQ ID NO:5).
5. An isolated GPRA polypeptide comprising an amino acid sequence that has at least 80% sequence identity to an amino acid sequence selected from the group consisting of SEQ. ID NOS: 3, 5, 7, 9, 11, 13 and 15 over a sequence comparison window of at least 40 amino acids when compared using the BLASTP algorithm with a wordlength (W) of 3, and the BLOSUM62 scoring matrix provided that the polypeptide includes a variant amino acid encoded by a variant form shown in Table 7.
6. An isolated GPRA polypeptide of claim 1 comprising the amino acid sequence of SEQ ID NO:3 provided that the sequence contains an amino acid substitution of Asn 107Ile, Arg241Ser, or Gln344Arg.
7. An isolated GPRA polypeptide of claim 1 comprising the amino acid sequence of SEQ ID NO:5 provided that the sequence contain as an amino acid substitution of Asn instead of Ile at codon position 107, Arg instead of Ser at codon position 241, and/or Thr instead of Ile at codon position 366.

8. An isolated nucleic acid encoding the GPRA polypeptide of any of the preceding claims.

9. The isolated nucleic acid of claim 8 that hybridizes under highly stringent conditions to any of SEQ ID NOS: 4, 6, 8, 10, 12, and 14 wherein the highly stringent conditions are $6 \times$ NaCl/sodium citrate (SSC) at about 45°C for a hybridization step, followed by a wash of $2 \times$ SSC at 50°C

10. An isolated nucleic acid of claim 8 that hybridizes under highly stringent conditions to any of SEQ ID NOS: 1, 4, 6, 8, 10, 12, and 14 without hybridizing under the same highly stringent conditions to SEQ ID NO:2, wherein the highly stringent conditions are $6 \times$ NaCl/sodium citrate (SSC) at about 45°C for a hybridization step, followed by a wash of $2 \times$ SSC at 50°C .

11. An isolated nucleic acid of claim 8 having a sequence that is at least 90 % identical to a nucleic acid having a sequence selected from the group consisting of SEQ ID NO: 4, 6, 8, 10, 12, and 14 over the entire length of the selected SEQ ID NO when compared using the BLASTN algorithm with a wordlength (W) of 11, $M=5$, and $N=-4$.

12. An isolated nucleic acid of claim 8 having a sequence that is at least 80 % identical to a nucleic acid having a sequence selected from the group consisting of SEQ ID NO: 1, 2, 4, 6, 8, 10, 12, and 14 over a sequence comparison window of at least 100 nucleotides when compared using the BLASTN algorithm with a wordlength (W) of 11, $M=5$, and $N=-4$ provided that the nucleic acid includes a polymorphic site occupied by a variant form as shown in Table 3 or Table 7.

13. The isolated nucleic acid of claim 12, wherein the variant form is at a polymorphic site not designated by a * in Table 7.

14. An isolated nucleic acid of claim 8 having a sequence that is at least 80 % identical to a nucleic acid having a sequence selected from the group consisting of SEQ ID NO: 1, 2, 4, 6, 8, 10, 12, and 14 over a sequence comparison window of at least 100 nucleotides when compared using the BLASTN algorithm with a wordlength (W) of 11, $M=5$, and $N=-4$ provided that the nucleic acid includes a polymorphic site occupied by a reference form designated with a * in Table 7.

15. An isolated genomic DNA molecule or a minigene having at least one intronic sequence and encoding a GPRA polypeptide that has at least 80 % sequence identity to an amino acid sequence selected from the group consisting of SEQ. ID NOS: 3, 5, 7, 9, 11, 13 and 15 over a region at least 40 amino acids in length when compared using the BLASTP algorithm with a wordlength (W) of 3, and the BLOSUM62 scoring matrix .
16. The isolated genomic DNA molecule or minigene of claim 15, comprising at least exons III-IV of SEQ ID NO: 2.
17. The isolated genomic DNA molecule or minigene of claim 15, wherein the intronic sequence is from intron 2 or 3 of a GPRA gene.
18. The isolated genomic DNA molecule or minigene of claim 15, wherein the GPRA polypeptide includes amino acids 343-377 of B-Long (SEQ ID NO:5)
19. The isolated genomic DNA molecule or minigene of claim 15, wherein the polypeptide is SEQ. ID NOS: 3, 5, 7, 9, 11, 13 or 15.
20. The isolated nucleic acid, genomic DNA molecule or minigene of any of claims 8-19, linked to a second nucleic acid with which it is not naturally associated.
21. The isolated nucleic acid, genomic DNA molecule or minigene of claim 20, wherein the second nucleic acid includes a heterologous promoter operably linked to a gene within the isolated nucleic acid.
22. A vector comprising the isolated nucleic acid, genomic molecule or minigene nucleic acid of any of claims 8-21.
23. A host cell comprising the vector of claim 22.
24. An antibody that specifically binds to an epitope within amino acids 343-377 of B-long (SEQ ID NO:5) or amino acids 332-366 of B-short (SEQ ID NO:7).
25. A method of preventing or treating asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer, comprising
administering to a patient suffering from or at risk of asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer an effective amount of a

modulator of a GPRA polypeptide comprising an amino acid sequence selected from the group consisting of SEQ. ID NOS: 3, 5, 7, 9, 11, 13, and 15.

26. The method of claim 25, further comprising administering an effective amount of a modulator of a AAA1 polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41.

27. The method of claim 25, wherein the modulator binds to the GPRA polypeptide.

28. The method of claim 25, wherein the modulator inhibits expression of the GPRA polypeptide.

29. The method of claim 25, wherein the modulator is a transcript of an AAA1 gene or an inhibitor of expression of an AAA1 gene.

30. The method of claim 26, wherein the modulator of the AAA1 polypeptide binds to the AAA1 polypeptide.

31. The method of claim 26, wherein the modulator of the AAA1 polypeptide inhibits expression of the AAA1 polypeptide.

32. A method of identifying a modulator of a GPRA polypeptide, comprising

contacting a cell expressing a GPRA polypeptide with an agent;
determining whether the agent modulates expression of the GPRA polypeptide and/or signal transduction through the GPRA polypeptide, wherein the GPRA polypeptide is defined by any of claims 1-7.

33. The method of claim 32, wherein the cell further expresses an AAA1 polypeptide.

34. A method of determining risk of asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer, comprising;

determining whether the individual has a variant polymorphic form in a GPRA gene, wherein presence of the variant polymorphic form indicates risk of asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer.

35. The method of claim 34, wherein the variant form occurs in a noncoding region of the GPRA gene

36. The method of claim 34, wherein the variant form occurs in a coding region of the GPRA gene

37. The method of claim 34, wherein the variant form occurs between introns 1 and 4 of the GPRA gene.

38. The method of claim 34, wherein the determining comprises determining whether the individual has a variant form relative to SEQ ID NO: 1 (AST-1 locus).

39. The method of claim 34, wherein the determining comprises determining whether the individual has a variant form relative to any of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, and 15.

40. The method of claim 34, wherein the variant form is a variant form shown in Table 3.

41. The method of claim 40, wherein the variant form is at a polymorphic site shown in Table 3.

42. The method of claim 34, wherein the variant form is a variant form shown in Table 7.

43. The method of claim 34, wherein the variant form is a variant form at a polymorphic site not designated * in Table 7.

44. The method of claim 34, wherein the determining comprising determining whether the individual has variant polymorphic forms relative to SEQ ID NO:1 at each of a plurality of polymorphic sites within the AST-1 locus, the presence of variant polymorphic forms at two or more of the plurality of polymorphic sites indicating increased risk of asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer.

45. The method of claim 34, further comprising determining whether the individual has a variant polymorphic form in an AAA1 gene, wherein presence of the variant

polymorphic form indicates risk of asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer.

46. The method of claim 45, wherein the variant polymorphic form occurs in the coding region of the AAA1 gene.

47. The method of claim 34, further comprising amplifying at least part of SEQ ID NO:1 (AST-1) locus including the polymorphic site before the determining step.

48. The method of claim 34, wherein the determining is performed by allele specific amplification, allele specific hybridization, single strand conformation polymorphism (SSCP), oligonucleotide ligation assay, single-base extension assay, or restriction fragment length polymorphism (RFLP).

49. A method for identifying a polymorphic site correlated with a disease selected from the group consisting of asthma, other IgE-mediated disease, chronic obstructive pulmonary disease and cancer or susceptibility thereto, comprising;

identifying a polymorphic site within a GPRA gene,

determining whether a variant polymorphic form occupying the site is associated with the disease or susceptibility thereto.

50. The method of claim 49, wherein the variant form occurs in a noncoding region of the GPRA gene

51. The method of claim 49, wherein the variant form occurs in a coding region of the GPRA gene

52. The method of claim 49, wherein the variant form occurs between introns 2 and 4 of the GPRA gene.

53. The method of claim 49, wherein the determining comprises determining whether the individual has a variant form relative to SEQ ID NO: 1 (AST-1 locus)

54. The method of claim 49, wherein the determining comprises determining whether the individual has a variant form relative to any of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, and 15.

55. The method of claim 49, wherein the determining is performed by comparing the frequency of the variant polymorphic form in individuals with and without the disease.

56. A primer or probe nucleic acid that hybridizes under highly stringent conditions to a segment of SEQ ID NO:1, 2 or 4 or a variant form thereof differing from SEQ ID NO: 1, 2 or 4 at a position shown in Table 3 or Table 7, wherein the segment includes or is immediately adjacent to a polymorphic site shown in Table 3 or Table 7.

57. The primer or probe of claim 56, wherein the position is a position other than a position designated * in Table 7.

58. The primer or probe of claim 56 that is perfectly complementary to a segment of SEQ ID NO:1, 2 or 4.

59. The primer or probe of claim 56 that is perfectly complementary to a variant form of a segment of SEQ ID NO:1, 2 or 4 shown in Table 3 or Table 7.

60. The primer or probe of claim 56 that specifically hybridizes to the segment of SEQ. ID NO: 1, 2 or 4 without hybridizing to a corresponding segment of an allelic variant shown in Table 3 or Table 7.

61. A primer of claim 56 for conducting a single-base extension reaction, whereby the primer is perfectly complementary to a segment that is immediately adjacent to but does not include the polymorphic site.

62. A transgenic animal comprising a nucleic acid according to any one of claims 8-21.

63. The transgenic animal of claim 62, further comprising a nucleic acid encoding an AAA1 polypeptide.

64. A transgenic animal of claim 62 disposed to develop a characteristic of asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer in

which an endogenous a GPRA gene encoding a cognate form of a GPRA polypeptide defined by any of SEQ ID NOS: 3, 5, 7, 9, 11, 13 and 15 is functionally disrupted to prevent expression of a gene product.

65. The transgenic animal of claim 64 in which an endogenous AAA1 gene encoding a cognate form of an AAA1 polypeptide defined by an of SEQ ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 is functionally disrupted to prevent expression of a gene product of the AAA1 gene.

66. A kit for use in diagnosing or assessing predisposition to asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer, comprising;

a container; and in the container:

a compound, preferably labeled, capable of detecting a polymorphic form at a polymorphic site in a susceptibility locus for asthma as defined by SEQ ID NO:1, 2 or 4.

67. The kit according to claim 66, wherein the polymorphic site occurs at a position shown in Table 3, Table 7, Table 12, Table 13 or Table 14.

68. The kit according to claim 66, wherein the compound is a primer or probe.

69. The kit according to claim 66, wherein said primer is the primer of claim 56.

70. An isolated AAA1 polypeptide comprising an amino acid sequence that has at least 80% sequence identity to an amino acid sequence selected from the group consisting of SEQ. ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 over the entire length of the selected SEQ ID No: when compared using the BLASTP algorithm with a wordlength (W) of 3, and the BLOSUM62 scoring matrix.

71. The isolated AAA1 polypeptide of claim 70 comprising an amino acid sequence that has at least 90% sequence identity to the selected amino acid sequence.

72. The isolated AAA1 polypeptide of claim 70 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41.

73. The isolated AAA1 polypeptide of claim 70 comprising at least 10 contiguous amino acids from an amino acid sequence selected from the group consisting of SEQ ID NOS:17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41.

74. An isolated nucleic acid encoding the AAA1 polypeptide of any of claims 70-73.

75. The isolated nucleic acid of claim 74 that hybridizes under highly stringent conditions to any of SEQ ID NOS: 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40, wherein the highly stringent conditions are $6 \times$ NaCl/sodium citrate (SSC) at about 45°C for a hybridization step, followed by a wash of $2 \times$ SSC at 50°C

76. An isolated nucleic acid of claim 74 having a sequence that is at least 80% identical to a nucleic acid having a sequence selected from the group consisting of SEQ ID NO: 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40 over the entire length of the selected SEQ ID NO when compared using the BLASTN algorithm with a wordlength (W) of 11, M=5, and N= -4.

77. The isolated nucleic acid of claim 76, wherein at least one polymorphic site shown in Table 12 is occupied by a variant nucleotide.

78. An isolated nucleic acid having at least 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOS:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40.

79. An isolated genomic DNA molecule or a minigene having at least one intronic sequence and encoding an AAA1 polypeptide that has at least at least 80 % sequence identity to an amino acid sequence selected from the group consisting of SEQ. ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 over the entire length of the selected SEQ ID NO. when compared using the BLASTP algorithm with a wordlength (W) of 3, and the BLOSUM62 scoring matrix .

80. The isolated genomic DNA molecule or minigene of claim 79, wherein the polypeptide has a sequence seleted from the group consisting of SEQ. ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41.

81. The isolated nucleic acid, genomic DNA molecule or minigene of claim 79 or 80 linked to a second nucleic acid with which it is not naturally associated.

82. The isolated nucleic acid, genomic DNA molecule or minigene of claim 79, wherein the second nucleic acid includes a heterologous promoter operably linked to a gene within the isolated nucleic acid.

83. A vector comprising the isolated nucleic acid, genomic molecule or minigene nucleic acid of any of claims 74-82.

84. A host cell comprising the vector of claim 83.

85. An antibody that specifically binds to a polypeptide selected from the group consisting of SEQ ID NOS:17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41.

86. A method of preventing or treating asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer, comprising

administering to a patient suffering from or at risk of asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer an effective amount of a modulator of an AAA1 polypeptide comprising an amino acid sequence selected from the group consisting of SEQ. ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41.

87. The method of claim 86, wherein the modulator binds to the AAA1 polypeptide.

88. The method of claim 86, wherein the modulator inhibits or induces expression of the AAA1 polypeptide.

89. A method of identifying a modulator of an AAA1 polypeptide, comprising

contacting an AAA1 polypeptide with an agent;
determining whether the agent binds to the AA1 polypeptide,
modulates expression of the AAA1 polypeptide or modulates activity of the AAA1 polypeptide, wherein the AAA1 polypeptide comprises an amino acid sequenced as defined by any of SEQ ID NOS:17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41.

90. The method of claim 89, wherein the AAA1 polypeptide is expressed from a cell.

91. A method of determining risk of asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer, comprising;

determining whether the individual has a variant polymorphic form in an AAA1 gene, wherein presence of the variant polymorphic form indicates risk of asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer.

92. The method of claim 91, wherein the variant form occurs in a noncoding region of the AAA1 gene

93. The method of claim 91, wherein the variant form occurs in a coding region of the AAA1 gene

94. The method of claim 91, wherein the determining comprises determining whether the individual has a variant form relative to SEQ ID NO: 1 (AST-1 locus)

95. The method of claim 91, wherein the determining comprises determining whether the individual has a variant form relative to any of SEQ ID NOS: 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40.

96. The method of claim 91, wherein the variant form is a variant form shown in Table 12.

97. The method of claim 91, wherein the determining is performed by allele specific amplification, allele specific hybridization, single strand conformation polymorphism (SSCP), oligonucleotide ligation assay, single-base extension assay, or restriction fragment length polymorphism (RFLP).

98. A method for identifying a polymorphic site correlated with a disease selected from the group consisting of asthma, other IgE mediated disease, chronic obstructive pulmonary disease or cancer or susceptibility thereto, comprising;

identifying a polymorphic site within an AAA1 gene,

determining whether a variant polymorphic form occupying the site is associated with the disease or susceptibility thereto.

99. The method of claim 98, wherein the variant form occurs in a noncoding region of the AAA1 gene.

100. The method of claim 98, wherein the variant form occurs in a coding region of the AAA1 gene

101. The method of claim 98, wherein the determining comprises determining whether the individual has a variant form relative to SEQ ID NO: 1 (AST-1 locus)

102. The method of claim 98, wherein the determining comprises determining whether the individual has a variant form relative to any of SEQ ID NOS: 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40.

103. The method of claim 98, wherein the determining is performed by comparing the frequency of the variant polymorphic form in individuals with and without the disease.

104. A primer or probe nucleic acid of nucleotides that hybridizes under highly stringent conditions to a segment of SEQ ID NO: 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40 or a variant form thereof differing from SEQ ID NOS.16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40 at a single polymorphic site.

105. The primer or probe of claim 104, wherein the polymorphic site is one shown in Table 3 or 12.

106. The primer or probe of claim 104 that is perfectly complementary to a segment of SEQ ID NOS:16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38 and 40.

107. A primer or probe of claim 104 for conducting a single-base extension reaction, whereby the primer is perfectly complementary to a segment that is immediately adjacent to but does not include the polymorphic site.

108. A transgenic animal comprising a nucleic acid according to any one of claims 74-82.

109. A transgenic animal of claim 108 disposed to develop a characteristic of asthma, other IgE-mediated disease, chronic obstructive pulmonary disease or cancer in

which an endogenous a AAA1 gene encoding a cognate form of an AAA1 polypeptide defined by any of SEQ ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39 and 41 is functionally disrupted to prevent expression of a gene product.

110. A pharmaceutical composition comprising an effective amount of a GPRA polypeptide comprising an amino acid sequence selected from the group consisting of SEQ. ID NOS: 3, 5, 7, 9, 11, 13, and 15 or a modulator thereof for the prevention or treatment of asthma or other IgE mediated disease, chronic obstructive pulmonary disease or cancer.

111. The pharmaceutical composition of claim 110, further comprising an effective amount of an AAA1 polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, and 41 or a modulator thereof.

112. The pharmaceutical composition of claim 110, wherein the modulator binds to the GPRA polypeptide and/or activates/inhibits expression of the GPRA polypeptide.

113. The pharmaceutical composition of claim 111, wherein the modulator is a transcript of an AAA1 gene or an activator/inhibitor of expression of an AAA1 gene.

114. The pharmaceutical composition of claim 111, wherein the modulator binds to the AAA1 polypeptide and/or activates/inhibits expression of the AAA1 polypeptide.

115. Use of a GPRA polypeptide comprising an amino acid sequence selected from the group consisting of SEQ. ID NOS: 3, 5, 7, 9, 11, and 13 or a modulator thereof for the manufacture of a medicament for the prevention or treatment of asthma or other IgE mediated disease, chronic obstructive pulmonary disease or cancer.

116. The use of claim 115, wherein a GPRA polypeptide comprising an amino acid sequence selected from the group consisting of SEQ. ID NOS: 3, 5, 7, 9, 11, 13, and 15 or a modulator thereof, and an AAA1 polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, and 41 or a modulator thereof are used for the manufacture of said medicament.

117. A pharmaceutical composition comprising an effective amount of an AAA1 polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, and 41 or a modulator thereof for the prevention or treatment of asthma or other IgE mediated disease, chronic obstructive pulmonary disease or cancer.

118. The pharmaceutical composition of claim 117, wherein the modulator binds to the AAA1 polypeptide and/or activates/inhibits expression of the AAA1 polypeptide.

119. Use of an AAA1 polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, and 41 or a modulator thereof for the manufacture of a medicament for the prevention or treatment of asthma or other IgE mediated disease, chronic obstructive pulmonary disease or cancer.